

Compactly Generated Domain Theory

Ingo Battenfeld, Matthias Schröder and Alex Simpson[†]

*LFCS, School of Informatics
University of Edinburgh, UK*

Received

We propose compactly generated monotone convergence spaces as a well-behaved topological generalization of directed-complete partial orders (dcpos). The category of such spaces enjoys the usual properties of categories of “predomains” in denotational semantics. Moreover, such properties are retained if one restricts to spaces with a countable pseudobase in the sense of E. Michael, a fact that permits connections to be made with computability theory, realizability semantics and recent work on the closure properties of topological quotients of countably based spaces (qcb spaces). We compare the standard domain-theoretic constructions of products and function spaces on dcpos with their compactly generated counterparts, showing that these agree in important cases though not in general.

1. Introduction

Domain theory was originally developed by Dana Scott in order to build mathematical models of recursion, datatypes and other programming language features. It has since blossomed into a rich mathematical theory, centred around the study of directed-complete partial orders (dcpos), and their Scott topologies, see e.g. (Abramsky and Jung 1994; Gierz *et al.* 2003) for overviews.

One would like domain theory to provide a flexible toolkit for modelling different aspects of computation. However, in spite of its many achievements, traditional dcpo-based domain theory has limitations in this regard. For example, Gordon Plotkin has pointed out that, although traditional domain theory can be used to model higher-order types (using cartesian-closed categories), computability for non-discrete datatypes (using ω -continuous dcpos), and computational effects (as free algebras for inequational theories), it is not capable of modelling all three in combination. Similarly, although one can build domain-theoretic models of the Girard-Reynolds’ polymorphic lambda-calculus, it is not known how to combine such models with computational effects, nor how to build models satisfying the parametricity properties that are vital for proving program equivalences.

[†] Research supported by an EPSRC research grant, “Topological Models of Computational Metalanguages”, and an EPSRC Advanced Research Fellowship (Simpson).

In order to address these weaknesses, it seems necessary to leave the familiar dcpo-based world of traditional domain theory. One possible alternative is to identify subcategories of domain-like structures within suitable “realizability” categories, see e.g. (Phoa 1990; Longley and Simpson 1997; Reus and Streicher 1999). However, while such categories do indeed resolve the problems identified above (though no comprehensive account of this has ever been published), important properties of traditional domain theory, such as the connections with topology (Gierz *et al.* 2003), are generally lost.

Fortunately, the situation is not always so bad. In (Simpson 2003), the third author observed that, in one particular realizability category built over Scott’s combinatory algebra $\mathcal{P}\omega$ (Scott 1976), a natural category of “predomains” (the *complete extensional objects*) can equivalently be presented as a category of topological spaces, called *topological predomains* in *op. cit.* The importance of this coincidence, a proof of which appears in the first author’s Diploma thesis (Battenfeld 2004), is that it opens up the possibility of obtaining the benefits of realizability-based notions of domain within an orthodox topological framework. Further, topological predomains extend the scope of traditional domain theory, by including familiar topological spaces such as the Euclidean reals and Cantor space, which make sense as datatypes, but whose topology is not the Scott topology on any underlying partial order.

The objective of the present paper is to provide a self-contained introduction to topological predomains aimed specifically at readers familiar with traditional domain theory and its topological connections. We thus ignore the realizability side of topological predomains entirely, for which the interested reader is referred to (Simpson 2003; Battenfeld 2004). Instead, we derive topological predomains from first principles, taking cartesian-closedness as our starting point.

For dcpo, cartesian-closedness is a consequence of the order-theoretic setting. Once more general topological spaces are allowed as predomains, more inclusive cartesian closed categories of topological spaces are required. There is thus a natural connection with the realm of so-called “convenient topology”, introduced by Ronnie Brown in (Brown 1963; Brown 1964) and popularized by Norman Steenrod (Steenrod 1967). Convenient topology is the study of categories of spaces enjoying additional useful properties, in particular cartesian closedness, that are not possessed by the category **Top** of all topological spaces. In (Brown 1963; Brown 1964; Steenrod 1967), the category of compactly generated Hausdorff spaces is shown to be one such cartesian closed category. Many other cartesian closed subcategories (and supercategories) of **Top**, have since been studied for similar reasons, see (Preuss 2002; Escardó *et al.* 2004) for recent overviews. The idea behind the present paper is to take such a cartesian closed category of topological spaces as the basis for developing a generalized domain theory.

In Section 3 we develop notions of predomain and domain within the cartesian closed category of compactly generated spaces (of course we do not restrict to Hausdorff spaces, because interesting domains are never Hausdorff). *Compactly generated predomains* are simply the compactly generated “monotone convergence spaces” in the sense of (Gierz *et al.* 2003), and *domains* are predomains with least element in the specialization order. The main results of the section establish that the compactly generated and monotone convergence space properties combine nicely with each other. Indeed we obtain cartesian

closed categories of predomains and domains that are exponential ideals of the category of compactly generated spaces, with the former subcategory being reflective.

In Section 4, we show that the good properties of compactly generated (pre)domains are retained under the imposition of a smallness condition on the topology. Here, the appropriate condition is to require a countable *pseudobase* in the sense of (Michael 1966), which generalizes the standard notion of base for a topology. In fact, the countably pseudobased compactly generated spaces have a simple characterization as the topological quotients of countably-based spaces (*qcb spaces*), which themselves form a cartesian closed category of topological spaces (Schröder 2003; Menni and Simpson 2002). In Section 4, we study the subcategory of monotone convergence qcb spaces, which coincides with the category of *topological predomains* introduced in (Simpson 2003), discussed above. We show that this category is a full reflective exponential ideal of the category of qcb spaces (this result was stated but not proved in *op. cit.*), hence it too is cartesian closed with countable limits and colimits.

One advantage of considering the larger category of *all* compactly generated predomains is that, by a result due to Jimmie Lawson (Escardó *et al.* 2004, Theorem 4.7), it contains the category of dcpos as a subcategory. Thus compactly generated predomains extend the world of traditional domain theory. In Section 5, we address the question of whether the traditional domain-theoretic constructions on dcpos, such as products and function spaces, agree with their compactly generated counterparts. Although products of domains always agree, function spaces differ in general. Nevertheless, in important cases where the domain-theoretic construction is known to be well behaved, we show that function spaces do coincide. In other cases, we suggest that it is the compactly generated topology that is the more reasonable choice.

The topic of this paper lies on the boundary between domain theory and general topology, two subjects that have enjoyed an extremely rich interaction ever since the inception of domain theory, see (Gierz *et al.* 2003) for a survey. It is a pleasure to dedicate this paper to Klaus Keimel, who has been one of the main contributors to the development of this interaction.

2. Preliminaries

Our notation and terminology is mainly standard. For a topological space X , we write $\mathcal{O}(X)$ for the family of open sets of X . For arbitrary (possibly non-Hausdorff) spaces we use *compact* to mean the Heine-Borel property. We write \mathbb{S} for Sierpinski space, which has underlying set $\{\perp, \top\}$ with $\{\top\}$ open but $\{\perp\}$ not.

In domain theory, we consider directed-complete partial orders (dcpos) and directed-complete pointed partial orders (dcppos), i.e. dcpos with least element. We use \sqsubseteq for the partial order structure in dcpos, \bigsqcup (resp. \bigsqcap) for suprema (resp. infima) and $\downarrow X$ (resp. $\uparrow X$) for the down-closure (resp. up-closure) of a subset in the order. For continuous dcpos we write \ll for the way-below relation, and use $\downarrow x$ (resp. $\uparrow x$) for the sets of elements way-below (resp. way-above) an element x .

3. Compactly generated predomains

We begin with an overview of compactly generated spaces, cf. (Brown 1963; Brown 1964; Steenrod 1967) (but without the Hausdorff restriction). We give full definitions, but state many standard properties without proof. For a recent comprehensive treatment see (Escardó *et al.* 2004).

Definition 3.1 (Compactly generated topology) A subset V of a topological space X is open in the *compactly generated topology* on X if, for every compact Hausdorff space K and continuous $p: K \rightarrow X$, the preimage $p^{-1}V$ is open in K . We write $k(X)$ for X with the compactly generated topology, and we say that X is *compactly generated* if $X = k(X)$.

Compactly generated spaces include all locally compact Hausdorff spaces. They also include a rich collection of non-Hausdorff spaces. For example, every sequential space is compactly generated. Hence, as special cases, all first-countable spaces are compactly generated, and so are ω -cpo's with the ω -Scott topology. A far less straightforward fact, due to Jimmie Lawson, is that compactly generated spaces also include all dcpos (with Scott topology). This result is pivotal to the development of this paper, so we state it as a proposition.

Proposition 3.2 *Every dcpo with its Scott topology is compactly generated.*

For a proof see Theorem 4.7 of (Escardó *et al.* 2004).

We write \mathbf{kTop} for the category of compactly generated spaces and continuous functions. As is well known (and easily seen), \mathbf{kTop} is a coreflective subcategory of \mathbf{Top} , where the coreflection functor maps X to $k(X)$. It follows that \mathbf{kTop} is cocomplete with colimits calculated as in \mathbf{Top} , and complete with limits obtained by coreflecting limits in \mathbf{Top} . In particular, the cartesian product in \mathbf{kTop} of a family $\{X_i\}_{i \in I}$ of compactly generated spaces is given by $k(\prod_{i \in I} X_i)$, where $\prod_{i \in I} X_i$ is the topological product. We write $\prod_{i \in I}^k X_i$ and $X \times_k Y$ for the products $k(\prod_{i \in I} X_i)$ and $k(X \times Y)$ in \mathbf{kTop} . In certain cases, this description of $X \times_k Y$ can be simplified. For example, if X and Y are countably based then so is $X \times Y$, and hence $X \times_k Y = X \times Y$. The proposition below gives another important case in which the topologies coincide. A topological space X is said to be *core compact* if its open sets form a continuous lattice under inclusion. Core compactness is a mild generalization of local compactness: every locally compact space is core compact, and every core compact sober space is locally compact. Core compact spaces arise naturally as the exponentiable objects in \mathbf{Top} .

Proposition 3.3 *If X, Y are compactly generated spaces and X is core compact then $X \times_k Y = X \times Y$.*

For a proof see Theorem 5.4 of (Escardó *et al.* 2004).

For topological spaces X, Y , we write $C(X, Y)$ for the set of continuous functions from X to Y .

Definition 3.4 (Compact open topology) For topological spaces X, Y , the *compact open topology* on $C(X, Y)$ is generated by the subbasic opens

$$\langle K, V \rangle = \{f \mid f(K) \subseteq V\} ,$$

where $K \subseteq X$ is compact and $V \subseteq Y$ is open. We write $C_{\text{co}}(X, Y)$ for $C(X, Y)$ with the compact open topology.

It is well known that every locally compact space X is exponentiable in **Top**, with the exponential Y^X given by $C_{\text{co}}(X, Y)$.

Proposition 3.5 *The category **kTop** is cartesian closed with exponential $X \Rightarrow_k Y$ given by $k(C_{\text{co}}(X, Y))$.*

This is a standard result, see (Escardó *et al.* 2004) for references and for a recent exposition of the general machinery underlying the construction of exponentials in cartesian closed subcategories of **Top**. (Although it does not appear explicitly in (Escardó *et al.* 2004), the coincidence of $X \Rightarrow_k Y$ and $k(C_{\text{co}}(X, Y))$ follows easily from Theorem 5.15 and Remark 5.20 of *op. cit.*, using the fact that the Isbell topology refines the compact open topology.)

The next two results give special cases in which the exponential in **kTop** is easily calculated.

Proposition 3.6 *If X, Y are countably based and X is locally compact then $C_{\text{co}}(X, Y)$ is countably based, hence $X \Rightarrow_k Y = C_{\text{co}}(X, Y)$.*

Proposition 3.7 *If X is compactly generated then $X \Rightarrow_k \mathbb{S}$ has the Scott topology.*

A proof of the first proposition can be found in (Lambrinos and Papadopoulos 1985). The second, which appears as Corollary 5.16 of (Escardó *et al.* 2004), is again due to Jimmie Lawson.

In traditional domain theory, the Scott topology is derived from the partial order. To define our notion of predomain, we also work with order-theoretic properties, but we take the topology as primary and the order as derived. Recall that the *specialization order* on a topological space is defined as follows.

Definition 3.8 (Specialization order) The *specialization order* \sqsubseteq on a topological space X is defined by $x \sqsubseteq y$ if, for all open $U \subseteq X$, $x \in U$ implies $y \in U$.

Trivially, every open set $U \subseteq X$ is upper-closed in the specialization order, which is, in general, a preorder on X . The space X is said to be T_0 if \sqsubseteq is a partial order.

Definition 3.9 (Monotone convergence space) A topological space X is a *monotone convergence space* if: the specialization order on X is a dcpo (in particular, X is T_0), and every open subset of X is Scott-open with respect to the order.

Monotone convergence spaces include: all T_1 spaces, all sober spaces, and all dcpos with the Scott topology. Monotone convergence spaces were introduced in (Wyler 1981), where they were called *d-spaces*. We take our terminology from (Gierz *et al.* 2003).

We now come to the central definition in this paper.

Definition 3.10 (Compactly generated predomain) A *compactly generated predomain* is a topological space X that is both compactly generated and a monotone convergence space.

We write \mathbf{kP} for the category of compactly generated predomains and continuous functions. By combining previous remarks, one sees that compactly generated predomains include all locally compact Hausdorff spaces and all dcpos with the Scott topology.

In order to state the main theorem of this section, recall that a full subcategory \mathcal{C}' of a cartesian-closed category \mathcal{C} is said to be an *exponential ideal* if it is closed under finite products and isomorphisms in \mathcal{C} and, for all objects X of \mathcal{C} and Y of \mathcal{C}' , the \mathcal{C} -exponential Y^X is an object of \mathcal{C}' . Obviously, exponential ideals are themselves cartesian closed.

Theorem 3.11 *The category \mathbf{kP} is a full reflective exponential ideal of \mathbf{kTop} .*

It follows that \mathbf{kP} is complete and cocomplete. Limits are calculated as in \mathbf{kTop} . Colimits are calculated by reflecting colimits from \mathbf{kTop} . Thus, in \mathbf{kP} , neither limits nor colimits are, in general, calculated as in \mathbf{Top} (though it is easy to see that coproducts in \mathbf{kP} are calculated as in \mathbf{Top}).

We prove the theorem in stages. First, we observe that the coreflection from \mathbf{Top} to \mathbf{kTop} cuts down to monotone convergence spaces.

Proposition 3.12 *If X is a monotone convergence space then so is $k(X)$.*

Proof. Since Sierpinski space \mathbb{S} is compactly generated, $C(\mathbb{S}, k(X)) = C(\mathbb{S}, X)$, thus $k(X)$ and X have the same specialization order, which is a dcpo. Easily, $k(X)$ is the coarsest compactly generated topology on the set X that refines the topology on X . Since X is a monotone convergence space, the Scott topology on the specialization order refines the topology on X . By proposition 3.2, the Scott topology is compactly generated. Hence every open in $k(X)$ is Scott open. \square

Lemma 3.13 *If X is a topological space and Y is a monotone convergence space then the pointwise order on $C(X, Y)$ is a dcpo with directed suprema constructed pointwise.*

For the straightforward proof see Lemma II.3.14 of (Gierz *et al.* 2003).

Proposition 3.14 *The category \mathbf{kP} is an exponential ideal of \mathbf{kTop} .*

Proof. For closure under finite products, one easily shows that topological products preserve monotone convergence spaces, from which the result follows by Proposition 3.12.

For the exponential property, suppose X is compactly generated and Y is a compactly generated predomain. By Propositions 3.5 and 3.12, it suffices to show that $C_{\text{co}}(X, Y)$

is a monotone convergence space. It is easily checked that the specialization order on $C_{\text{co}}(X, Y)$ is pointwise, and this is a dcpo by Lemma 3.13. It remains to show that every subbasic open $\langle K, V \rangle$ of $C_{\text{co}}(X, Y)$ is Scott open. But suppose $(\bigsqcup_{i \in I} f_i) \in \langle K, V \rangle$, where $\{f_i\}_{i \in I}$ is directed. Because V is Scott open, $\{f_i^{-1}V\}_{i \in I}$ is an open cover of K . By directedness, there exists $j \in I$ such that f_j is an upper bound for finitely many f_i determining a finite subcover. Then $f_j \in \langle K, V \rangle$. Thus $\langle K, V \rangle$ is indeed Scott open. \square

One way of obtaining the reflection functor from \mathbf{kTop} to \mathbf{kP} would be to apply the Special Adjoint Functor Theorem. Instead, we provide a more informative direct construction. We show that the reflection from \mathbf{Top} to the category of monotone convergence spaces, as described in (Wyler 1981), cuts down to a reflection from \mathbf{kTop} to \mathbf{kP} . This nicely mirrors the symmetric property established in Proposition 3.12 for the coreflection k .

First, we present the reflection from \mathbf{Top} to the category of monotone convergence spaces, cf. (Wyler 1981). Recall that a filter $\mathcal{F} \subseteq \mathcal{O}(X)$ is said to be *completely prime* if whenever $(\bigcup_{i \in I} U_i) \in \mathcal{F}$ it holds that $U_i \in \mathcal{F}$ for some $i \in I$. (This implies that $\emptyset \notin \mathcal{F}$.) For any point $x \in X$, the filter $\eta(x)$ of open neighbourhoods of x is always completely prime. A topological space is said to be *sober* if every completely prime filter is the filter of open neighbourhoods of a unique point. The *sobrification* $\mathfrak{S}(X)$ of a topological space X has the set of completely prime filters of $\mathcal{O}(X)$ as its underlying set with open sets

$$\{\mathcal{F} \in \mathfrak{S}(X) \mid U \in \mathcal{F}\},$$

where $U \in \mathcal{O}(X)$. One easily observes that specialization order on $\mathfrak{S}(X)$ is inclusion, with least upper bounds of directed subsets $D \subseteq \mathfrak{S}(X)$ given by $\bigcup D$, which is indeed a completely prime filter. Define $\mathfrak{M}(X)$ to be the smallest subspace of $\mathfrak{S}(X)$ that contains all neighbourhood filters and is closed under directed lubs in the specialization order. It will be useful to have an explicit description of the topology on $\mathfrak{M}(X)$.

Lemma 3.15 *The following are equivalent for a subset $V \subseteq \mathfrak{M}(X)$.*

- 1 V is open.
- 2 $\eta^{-1}V$ is an open subset of X and $V = \uparrow(V \cap \eta(X))$ in the specialization order on $\mathfrak{M}(X)$.
- 3 $\eta^{-1}V$ is an open subset of X and V is Scott-open in the specialization order on $\mathfrak{M}(X)$.

Proof. For 2 \implies 1, suppose that $\eta^{-1}V$ is open in X and $V = \uparrow(V \cap \eta(X))$. We show that $V = \{\mathcal{F} \in \mathfrak{M}(X) \mid (\eta^{-1}V) \in \mathcal{F}\}$, and hence V is open by the definition of the topology on $\mathfrak{S}(X)$. If $\mathcal{F} \supseteq \eta(x)$ for $x \in \eta^{-1}V$ then trivially $(\eta^{-1}V) \in \mathcal{F}$. Thus $\{\mathcal{F} \in \mathfrak{M}(X) \mid (\eta^{-1}V) \in \mathcal{F}\} \supseteq \uparrow(V \cap \eta(X)) = V$. Conversely, suppose that $(\eta^{-1}V) \in \mathcal{F} \in \mathfrak{M}(X)$. For any $x \in \eta^{-1}V$ for which $\eta(x) \not\subseteq \mathcal{F}$, there exists open $U_x \ni x$ in X such that $U_x \notin \mathcal{F}$. Suppose, for contradiction, that such U_x exists for every $x \in \eta^{-1}V$. Then $\bigcup_{x \in \eta^{-1}V} U_x \supseteq \eta^{-1}V$, so $\bigcup_{x \in \eta^{-1}V} U_x \in \mathcal{F}$. Hence, because \mathcal{F} is completely prime, $U_x \in \mathcal{F}$ for some $x \in \eta^{-1}V$, a contradiction. Thus there exists $x \in \eta^{-1}V$ with $\eta(x) \subseteq \mathcal{F}$. So indeed $\{\mathcal{F} \in \mathfrak{M}(X) \mid (\eta^{-1}V) \in \mathcal{F}\} \subseteq \uparrow(V \cap \eta(X)) = V$.

For 1 \implies 3, suppose $V \subseteq \mathfrak{M}(X)$ is open, i.e. there exists open $U \subseteq X$ such that $V = \{\mathcal{F} \in \mathfrak{M}(X) \mid U \in \mathcal{F}\}$. Then $\eta^{-1}V = U$, so $\eta^{-1}V$ is indeed an open subset of

X . Also, V is obviously upwards closed in the specialization order. To show that V is inaccessible by directed suprema, suppose that $D \subseteq \mathfrak{M}(X)$ is directed with $\bigsqcup D \in V$, i.e. $\bigcup D \in V$. Then $U \in \bigcup D$, so there exists $\mathcal{F} \in D$ with $U \in \mathcal{F}$, whence $\mathcal{F} \in V$ as required.

For $3 \implies 2$, suppose that $\eta^{-1}V$ is open in X and V is Scott-open in $\mathfrak{M}(X)$. Obviously $V \supseteq \uparrow(V \cap \eta(X))$, because Scott-open sets are upper closed. It remains to show that $V \subseteq \uparrow(V \cap \eta(X))$. Because V is Scott-open, it is contained in the Scott-interior of $\mathfrak{M}(X) \setminus \eta(X \setminus \eta^{-1}V)$. It is enough to show that the Scott-interior of $\mathfrak{M}(X) \setminus \eta(X \setminus \eta^{-1}V)$ is $\uparrow(V \cap \eta(X))$. We prove the equivalent statement that the Scott-closure of $\eta(X \setminus \eta^{-1}V)$ is $\mathfrak{M}(X) \setminus \uparrow(V \cap \eta(X))$. Let S be the Scott-closure of $\eta(X \setminus \eta^{-1}V)$. Then

$$S \cup \uparrow(V \cap \eta(X)) = \mathfrak{M}(X) , \quad (1)$$

because the left-hand side contains $\eta(X)$ and is closed under suprema of directed sets. Also, $S \cap V = \emptyset$ because $\eta(X \setminus \eta^{-1}V) \cap V = \emptyset$ and V is Scott-open. Hence

$$S \cap \uparrow(V \cap \eta(X)) = \emptyset , \quad (2)$$

because $\uparrow(V \cap \eta(X)) \subseteq V$. So, by (1) and (2), $S = \mathfrak{M}(X) \setminus \uparrow(V \cap \eta(X))$ as required. \square

We remark that the equivalence of 1 and 2 above is inherited by $\mathfrak{M}(X)$ from an analogous characterization of open sets in $\mathfrak{S}(X)$. That property 3 characterizes open sets is, however, a feature specific to $\mathfrak{M}(X)$.

It follows from Lemma 3.15 that $\mathfrak{M}(X)$ is a monotone convergence space. In fact it is the free monotone convergence space over X :

Proposition 3.16 *For any topological space X , the space $\mathfrak{M}(X)$ is a monotone convergence space. Moreover, for any monotone convergence space Y and continuous function $f: X \rightarrow Y$ there exists a unique continuous $g: \mathfrak{M}(X) \rightarrow Y$ such that $g \circ \eta = f$.*

Proof. This is Theorem 2.7 of (Wyler 1981). \square

Proposition 3.17 *If X is compactly generated then so is $\mathfrak{M}(X)$.*

Proof. Suppose $V \subseteq \mathfrak{M}(X)$ is such that for every compact Hausdorff K and continuous $p: K \rightarrow \mathfrak{M}(X)$ it holds that $p^{-1}V$ is open in K . We use Lemma 3.15 to show that V is open in $\mathfrak{M}(X)$, establishing condition 3. To show that $\eta^{-1}V$ is open in X , suppose that K is compact Hausdorff and $q: K \rightarrow X$ is continuous. Then $q^{-1}(\eta^{-1}V) = (\eta \circ q)^{-1}V$, so $q^{-1}(\eta^{-1}V)$ is open in K . Thus indeed $\eta^{-1}V$ is open in X , because X is compactly generated. It remains to show that V is Scott-open in $\mathfrak{M}(X)$. By assumption, V is open in $k(\mathfrak{M}(X))$, which has the same specialization order as $\mathfrak{M}(X)$, and is a monotone convergence space by Proposition 3.12. Thus indeed V is Scott-open in $\mathfrak{M}(X)$. \square

In combination, Propositions 3.14, 3.16 and 3.17 prove Theorem 3.11.

In domain theory, one is often interested in domains (i.e. dcpos) as opposed to pre-domains (i.e. dcpos). We make the analogous definition for compactly generated spaces.

Definition 3.18 (Compactly generated domain) *A compactly generated domain is a compactly generated pre-domain with a least element in the specialization order.*

We write \mathbf{kD} for the category of compactly generated domains and continuous functions.

Proposition 3.19 *The category \mathbf{kD} is an exponential ideal of \mathbf{kTop} , closed under arbitrary products.*

Proof. Given Theorem 3.11, all that remains to check is that the required constructions preserve the property of having a least element. This is straightforward. \square

The category of compactly generated domains is a category, properly extending the category of dcpops, that enjoys all the usual properties of a category of domains. Indeed, one can show that the category \mathbf{kD} enjoys the usual interrelationship with its subcategory of *strict* (i.e. bottom-preserving) maps, that the expected strict type constructors (smash product, strict function space, coalesced sum) are available, that recursive domain equations have solutions, etc. The constructions, which are routine modifications of the familiar domain-theoretic ones, are omitted from the present paper.

4. Countably pseudobased spaces

In traditional domain theory, countable (domain-theoretic) bases allow a theory of computability for domains to be developed. Such bases exist for all ω -continuous dcpops. Although the categories of ω -continuous dcpops and dcpos are not cartesian closed, they have cartesian-closed subcategories which, for many purposes (modulo the limitations discussed in Section 1), do provide workable categories of (pre)domains.

In our more general topological setting, a natural first attempt at doing something similar is to restrict to compactly generated spaces with countable (topological) bases. As with the category of ω -continuous dcpops, this category is not cartesian closed. In this case, it seems that the most natural remedy is to enlarge the category rather than to shrink it. This is done by weakening the requirement of a countable base to a countable *pseudobase* in the sense of E. Michael (Michael 1966).

Definition 4.1 (Pseudobase) A *pseudobase* for a topological space X is a family \mathcal{B} of (not necessarily open) subsets of X satisfying, whenever $K \subseteq U$ with K compact and U open subsets of X , there exist finitely many $B_1, \dots, B_k \in \mathcal{B}$ such that $K \subseteq \bigcup_{i=1}^k B_i \subseteq U$. A *pseudosubbase* is a family of subsets whose closure under finite intersection forms a pseudobase.

Obviously any (sub)base for the topology on X is also a pseudo(sub)base. Conversely, whenever a pseudo(sub)base \mathcal{B} consists of open sets then it is itself a (sub)base.

The requirements on a pseudobase are weak enough that it need have very little to do with the topology. For example, the powerset of X is always a pseudobase for X . However, as the results below demonstrate, pseudobases do become interesting when cardinality restrictions are placed upon them.

We say that a topological space X is a *qcb* (*quotient of countably based*) *space*, if it can be exhibited as a topological quotient $q: A \twoheadrightarrow X$, where A is a countably based space. We write \mathbf{QCB} for the full subcategory of \mathbf{Top} consisting of such spaces. In his PhD

thesis, the second author established that qcb spaces are exactly the sequential spaces with countable pseudobase (Schröder 2003). The proposition below, which follows from Theorem 6.10 of (Escardó *et al.* 2004), generalises this result to compactly generated spaces.

Proposition 4.2 *The following are equivalent for a topological space X .*

- 1 X is compactly generated and has a countable pseudobase.
- 2 X is a qcb space.

(To obtain this result as consequence of Theorem 6.10 of (Escardó *et al.* 2004), let \mathcal{C} be the category of compact Hausdorff spaces. Then every \llcorner -pseudobase in the sense of *loc. cit.* is a pseudobase as defined above, and every pseudobase as defined above is a $\llcorner_{\mathcal{C}}$ -pseudosubbase in the sense of *loc. cit.*)

Trivially, every countably based space is itself a qcb space. However, not every qcb space is countably based. The next result gives a useful sufficient condition under which qcb spaces are countably based.

Proposition 4.3 *If a locally compact space has a countable pseudobase then it has a countable base.*

Proof. The interiors of pseudobase sets form a base when the topology is locally compact, cf. Corollary 6.11 of (Escardó *et al.* 2004). \square

The next proposition reviews some of the useful properties of countably based spaces that are shared by the more general class of qcb spaces.

Proposition 4.4 *If X is a qcb space then:*

- 1 X is a sequential space.
- 2 X is hereditarily Lindelöf (i.e. for every family of opens $\{U_i\}_{i \in I}$ there exists countable $J \subseteq I$ such that $\bigcup_{i \in I} U_i = \bigcup_{j \in J} U_j$).
- 3 X is hereditarily separable (i.e. for any $Y \subseteq X$ there exists a countable $Y' \subseteq Y$ such that Y' is dense in the subspace topology on Y).

Proof. Properties 1 and 2 hold of countably based spaces and are preserved under quotienting. For 3, every space with countable pseudobase is separable, and pseudobases restrict to subspaces. (Note that the subspace topology on Y need not itself be compactly generated.) \square

Quite unexpectedly, the category **QCB** has very rich structure.

Proposition 4.5 *The category **QCB** has all countable limits and colimits and is cartesian closed. Moreover, this structure is preserved by the inclusion $\mathbf{QCB} \hookrightarrow \mathbf{kTop}$.*

This result is a special case of Corollary 7.3 of (Escardó *et al.* 2004), where a full proof is given. Earlier proofs of cartesian closedness appear in (Schröder 2003; Menni and

Simpson 2002). Here we simply state that if \mathcal{A}, \mathcal{B} are countable pseudosubbases for qcb spaces X, Y respectively, then the family of all sets of the form

$$\{f \in C(X, Y) \mid f(\bigcap \mathcal{A}') \subseteq \bigcup \mathcal{B}'\} ,$$

where \mathcal{A}' is a finite subset of \mathcal{A} and \mathcal{B}' is a finite subset of \mathcal{B} , form a countable pseudo-subbase for $X \Rightarrow_k Y$.

The goal of this section is to show that countably pseudobased compactly generated spaces (i.e. qcb spaces) form a good environment for restricting the notions of predomain and domain from the previous sections. We write $\omega\mathbf{P}$ and $\omega\mathbf{D}$ for the full subcategories of \mathbf{kP} and \mathbf{kD} respectively whose objects are qcb spaces. Clearly $\omega\mathbf{P}$ (resp. $\omega\mathbf{D}$) contains every ω -continuous dcpo (resp. dcppo).

As is well known, ω -continuous dcpos can be equivalently defined using ω -completeness rather than directed-completeness. The proposition below shows that countable pseudobases permit an analogous flexibility in the definition of compactly generated predomain.

As is standard, we call an ascending sequence $x_0 \sqsubseteq x_1 \sqsubseteq x_2 \sqsubseteq \dots$ in a partial order an ω -chain. An ω -complete partial order (ω -cpo) is a partial order in which every ω -chain has a lub. A subset X of an ω -cpo D is said to be ω -Scott open if it is upper closed and, whenever $(\bigsqcup_i x_i) \in X$, for an ω -chain (x_i) , it holds that $x_i \in X$ for some i .

Definition 4.6 (Monotone ω -convergence space) A topological space X is a *monotone ω -convergence space* if: the specialization order on X is an ω -complete partial order, and every open subset of X is ω -Scott-open with respect to the order.

Proposition 4.7 *A qcb space is a monotone convergence space if and only if it is a monotone ω -convergence space.*

Proof. It is immediate that any space that is a monotone convergence space is a monotone ω -convergence space. For the converse, suppose that X is a monotone ω -convergence qcb space. To show that the specialization order is a dcpo, suppose $D \subseteq X$ is directed. We must show that $\bigsqcup D$ exists. By Proposition 4.4, X is hereditarily separable, so D considered as a subspace of X has a countable dense subset $\{d_i \mid i \in \mathbb{N}\} \subseteq D$. Because D is directed, we can construct $\{e_i \mid i \in \mathbb{N}\} \subseteq D$ such that each e_i is an upper bound for the finite set $\{d_i\} \cup \{e_j \mid j < i\}$. Obviously $e_0 \sqsubseteq e_1 \sqsubseteq e_2 \dots$ is an ascending sequence. Define $e_\infty = \bigsqcup_i e_i$. We claim that $e_\infty = \bigsqcup D$. To see it is an upper bound, suppose $d \in D$. To show that $d \sqsubseteq e_\infty$, suppose that $d \in U \subseteq X$ where U is open. We must show that $e_\infty \in U$. Because $\{d_i \mid i \in \mathbb{N}\} \subseteq D$ is dense, there exists $d_i \in U$. Hence indeed $e_\infty \in U$, because $d_i \sqsubseteq e_i \sqsubseteq e_\infty$. For leastness, suppose e is any upper bound for D . To show that $e_\infty \sqsubseteq e$, suppose $e_\infty \in U \subseteq X$ where U is open. Because X is an ω -convergence space there exists i such that $e_i \in U$. But $e_i \sqsubseteq e$ because $e_i \in D$. So indeed $e \in U$.

It remains to show that every open is Scott-open. Suppose $U \subseteq X$ is open $D \subseteq X$ is directed $\bigsqcup D \in U$. We must show that $d \in U$ for some $d \in D$. But, as above, $\bigsqcup D = \bigsqcup_i e_i$ hence $e_i \in U$ for some i . thus $d = e_i$ is the required element of D . \square

Theorem 4.8 *The category $\omega\mathbf{P}$ is a full reflective exponential ideal of \mathbf{QCB} .*

It follows that $\omega\mathbf{P}$ is cartesian closed with countable limits and colimits, where limits are calculated as in **QCB**.

The proof of Theorem 4.8 follows similar lines to that of the analogous Theorem 3.11.

Proposition 4.9 *The category $\omega\mathbf{P}$ is an exponential ideal of **QCB**.*

Proof. Immediate from Propositions 3.14 and 4.5. \square

To establish the reflection part of Theorem 4.8, we show that the reflection \mathfrak{M} of monotone convergence spaces in **Top** cuts down to **QCB**.

Proposition 4.10 *If X is a qcb space then so is $\mathfrak{M}(X)$.*

Proof. We first make the following observations. If Y is compactly generated then, by the cartesian closedness of **kTop**, the function $i: Y \rightarrow ((Y \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S})$, defined by $i(y) = \lambda U. U(y)$, is continuous. Also by cartesian closedness, for any open $U \subseteq Y$, the function $p_U: ((Y \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S}) \rightarrow \mathbb{S}$, defined by $p_U(F) = F(U)$ (here we are identifying U with its characteristic function in $Y \Rightarrow_k \mathbb{S}$), is continuous. An easy calculation shows that, for any $y \in Y$ and open $U \subseteq Y$, it holds that $i(y) \in p_U^{-1}\{\top\}$ iff $y \in U$. Thus $i: Y \rightarrow ((Y \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S})$ is a topological pre-embedding. Further, if Y is T_0 then i is an injective function, and hence an embedding.

To prove the proposition, suppose that X is a qcb space. By Proposition 3.17, $\mathfrak{M}(X)$ is compactly generated. We must show that it also has countable pseudobase. By the observations above, there is a topological embedding of $\mathfrak{M}(X)$ in $((\mathfrak{M}(X) \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S})$. By Lemma 3.15, the function mapping $V \in \mathcal{O}(\mathfrak{M}(X))$ to $\eta^{-1}V$ gives a lattice isomorphism $\mathcal{O}(\mathfrak{M}(X)) \cong \mathcal{O}(X)$, whence, by Proposition 3.7, there is an induced homeomorphism $(\mathfrak{M}(X) \Rightarrow_k \mathbb{S}) \cong (X \Rightarrow_k \mathbb{S})$, and hence $((\mathfrak{M}(X) \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S}) \cong ((X \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S})$. Thus there is a topological embedding of $\mathfrak{M}(X)$ in $(X \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S}$. However, $(X \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S}$ is a qcb space because, by Theorem 4.8, qcb spaces are closed under function spaces in **kTop**. Thus $(X \Rightarrow_k \mathbb{S}) \Rightarrow_k \mathbb{S}$ has a countable pseudobase, and hence $\mathfrak{M}(X)$ does too, since pseudobases restrict to subspaces. \square

Theorem 4.8 now follows from Propositions 4.9, 3.16 and 4.10.

Proposition 4.11 *The category $\omega\mathbf{D}$ is an exponential ideal of **QCB**, closed under countable products.*

Proof. This is immediate from Propositions 3.19, 4.5 and Theorem 4.8. \square

The categories $\omega\mathbf{P}$ and $\omega\mathbf{D}$ were introduced in (Simpson 2003), where their objects were called *topological predomains* and *topological domains* respectively. The countable pseudobase requirement is sufficient for developing a computability theory, due to the connections, established in the second author's PhD thesis (Schröder 2003), between qcb spaces and Klaus Weihrauch's theory of *type two effectivity* (Weihrauch 2000). As outlined in (Simpson 2003), the categories of topological predomains and domains support the usual constructions of traditional domain theory, and also overcome the limitations discussed in Section 1. The details of this will appear elsewhere.

5. Comparison with traditional domain theory

Traditional domain theory concerns the categories **dcpo** and **dcppo** of continuous functions between dcpos and dcpos, and subcategories of them. By Proposition 3.2, **dcpo** and **dcppo** are full subcategories of **kP** and **kD** respectively. As is well known, **dcpo** and **dcppo** are themselves cartesian closed. Products $\prod_{i \in I}^s D_i$ and $D \times_s E$ of $\text{dc}(\text{p})\text{pos}$ are given by the product partial order. The exponential $D \Rightarrow_s E$ is given by the set of Scott continuous functions ordered pointwise.

In this section, we investigate the extent to which the cartesian closed structures on **dcpo** and **kP** agree. For finite products, there is no difference.

Proposition 5.1 *For dcpos D, E , it holds that $D \times_s E = D \times_k E$.*

Proof. Let D, E be dcpos. Then:

$$\begin{aligned} \mathcal{O}(D \times_s E) &\cong C(D \times_s E, \mathbb{S}) \\ &\cong C(D, E \Rightarrow_s \mathbb{S}) && \text{cartesian closedness of } \mathbf{dcpo} \\ &\cong C(D, E \Rightarrow_k \mathbb{S}) && \text{by Proposition 3.7} \\ &\cong C(D \times_k E, \mathbb{S}) && \text{cartesian closedness of } \mathbf{kP} \\ &\cong \mathcal{O}(D \times_k E) . \end{aligned}$$

So $D \times_s E$ and $D \times_k E$ carry the same topology. □

We remark that Martín Escardó has independently obtained the same result (Escardó 2005).

The above proposition shows that the inclusion **dcpo** \hookrightarrow **kP** preserves finite products. It does not preserve infinite products. For example the countable power of the two point discrete space is discrete in **dcpo**, but has the topology of Cantor space in **kP**. This counterexample makes essential use of a non-pointed space.

Proposition 5.2 *For any family $\{D_i\}_{i \in I}$ of dcpos, the Scott product $\prod_{i \in I}^s D_i$ and compactly generated product $\prod_{i \in I}^k D_i$ coincide.*

Proof. It is obvious that $\prod_{i \in I}^s D_i$ refines $\prod_{i \in I}^k D_i$, so we establish the converse. For any finite $J \subseteq I$, consider the set-theoretic function $\rho_J: \prod_{i \in I} D_i \rightarrow \prod_{i \in I} D_i$ defined by

$$(\rho_J(\pi))i = \begin{cases} \pi_i & \text{if } i \in J \\ \perp_{D_j} & \text{otherwise} \end{cases} .$$

Using the universal property of products in **kTop** and **dcppo** respectively, the two functions $\rho_J: \prod_{i \in I}^k D_i \rightarrow \prod_{i \in I}^k D_i$ and $\rho_J: \prod_{i \in I}^s D_i \rightarrow \prod_{i \in I}^s D_i$ are continuous idempotents, whose splittings are the retracts:

$$\begin{array}{ccc} \prod_{i \in I}^k D_i & \xrightarrow{r_k} & \prod_{j \in J}^k D_j & \xrightarrow{s_k} & \prod_{i \in I}^k D_i \\ \prod_{i \in I}^s D_i & \xrightarrow{r_s} & \prod_{j \in J}^s D_j & \xrightarrow{s_s} & \prod_{i \in I}^s D_i \end{array}$$




Fig. 1.

By Proposition 5.1, the finite products $\prod_{j \in J}^k D_j$ and $\prod_{j \in J}^s D_j$ are homeomorphic, thus the function $\rho_J = s_s \circ r_k: \prod_{i \in I}^k D_i \rightarrow \prod_{i \in I}^s D_i$ is continuous. Hence $\{\rho_J\}_J$ forms a directed family of continuous functions from $\prod_{i \in I}^k D_i$ to $\prod_{i \in I}^s D_i$, indexed by finite subsets $J \subseteq I$. Clearly, the pointwise supremum of this family is the identity function from $\prod_{i \in I}^k D_i$ to $\prod_{i \in I}^s D_i$. By Lemma 3.13, this is continuous, as required. \square

The above argument is adapted from Reinhold Heckmann’s proof of the analogous Theorem 7.8 in (Heckmann 2003).

The countable power of the two point discrete space again demonstrates a disagreement between function spaces in **dcpo** and **kP**: the function space $\mathbb{N} \Rightarrow_s \{0, 1\}$ is discrete, whereas $\mathbb{N} \Rightarrow_k \{0, 1\}$ is Cantor space. However, traditional domain theory largely concerns pointed dcpos. In the remainder of this section, we investigate the relationship between function spaces in **dcppo** and in **kD**. We shall see that the inclusion **dcppo** \hookrightarrow **kD** does not always preserve function spaces, but it does preserve them in many interesting cases. In fact, our claim is that the inclusion preserves function spaces in exactly those cases in which dcpo function spaces are “well behaved”. In other cases, **kD** defines a more reasonable function space than **dcppo**.

We begin with a counterexample to show that function spaces are not in general preserved by the inclusion **dcppo** \hookrightarrow **kD**. Consider the two (ω -algebraic) dcpos L_1 and L_2 presented in Fig. 1.

Proposition 5.3 *The function space $L_1 \Rightarrow_k L_2$ in **kP** does not have the Scott topology.*

Proof. Both L_1 and L_2 are countably based spaces, hence $L_1 \Rightarrow_k L_2$ is a qcb space. The Scott topology on the function space (i.e. $L_1 \Rightarrow_s L_2$) has been calculated by Achim Jung (Jung 1989). The resulting dcpo is algebraic, but not countably based (it has 2^{\aleph_0} compact elements). Because it is algebraic, the topology on $L_1 \Rightarrow_s L_2$ is locally compact. But then $L_1 \Rightarrow_s L_2$ cannot be a qcb space, for this would contradict Proposition 4.3. \square

An identical argument shows that the function space $L_1 \Rightarrow_k L_1$ in **kP** does not carry the Scott topology. The reason for choosing L_2 above was to give a counterexample with finite poset as codomain.

The dcpos L_1 and L_2 are both algebraic L-domains in the sense of (Jung 1989) (i.e. algebraic dcpos in which every principal ideal is a complete lattice). It is shown

in *op. cit.* that the category of algebraic (resp. continuous) L-domains forms one of the two maximal cartesian-closed categories of algebraic (resp. continuous) dcpos. One might thus be tempted to think of L-domains as belonging to the “well-behaved” part of traditional domain theory. But this disregards the fact that the ω -algebraic and ω -continuous L-domains do not form cartesian closed categories, due to the loss of countable base in the construction of function spaces. For the sake of contrast, by Proposition 3.6, the function space $L_1 \Rightarrow_k L_2$ in \mathbf{kD} is $C_{\text{co}}(L_1, L_2)$, which is countably based. In our view, it is the compactly generated function space that is the better behaved of the two.

When D is a continuous dcpo, the compact open topology $C_{\text{co}}(D, Y)$ can be given a simpler description.

Definition 5.4 (Point open topology) For topological spaces X, Y , the *point open topology* on $C(X, Y)$ is generated by the subbasic opens

$$\langle x, V \rangle = \{f \mid f(x) \subseteq V\} ,$$

where $x \in X$ and $V \subseteq Y$ is open. We write $C_{\text{po}}(X, Y)$ for $C(X, Y)$ with the compact open topology.

The point open topology is equivalently characterized as the topology of pointwise convergence, or as the relative topology on $C(X, Y)$ as a subspace of the product topology on the power Y^X . Trivially, the compact open topology $C_{\text{co}}(X, Y)$ always refines the point open topology $C_{\text{po}}(X, Y)$. When X is a continuous dcpo, the two agree.

Lemma 5.5 *If D is a continuous dcpo and Y a topological space then $C_{\text{po}}(D, Y)$ and $C_{\text{co}}(D, Y)$ coincide.*

Proof. Suppose $\langle K, V \rangle$ is a subbasic compact open neighbourhood of f . Then $K \subseteq f^{-1}(V)$ and so, as D is a continuous dcpo, $K \subseteq \bigcup_{x \in f^{-1}(V)} \hat{\uparrow}x$. As K is compact, there exists a finite $F \subseteq f^{-1}(V)$ such that $K \subseteq \bigcup_{x \in F} \hat{\uparrow}x$. But now we have $\bigcap_{x \in F} \langle x, V \rangle \subseteq \langle K, V \rangle$, and therefore $f \in \bigcap_{x \in F} \langle x, V \rangle \subseteq \langle K, V \rangle$, showing that $\langle K, V \rangle$ is point open, as required. \square

Together with Proposition 3.6, the above lemma, which is part of domain-theoretic folklore, implies that for ω -continuous dcpos D, E , we always have $D \Rightarrow_k E = C_{\text{co}}(D, E) = C_{\text{po}}(D, E)$, and this is countably based.

We have seen that $D \Rightarrow_k E$ does not always carry the Scott topology for ω -continuous D, E , even when D, E are L-domains. We next switch attention to the other of the two maximal cartesian-closed categories of dcpos, the category of FS-domains, introduced by Achim Jung (Jung 1990).

Definition 5.6 (FS-domain) An *FS-domain* is a dcpo D for which there exists a directed family $(f_i)_{i \in I}$ of continuous endofunctions on D , each *strongly finitely separated* from id_D , i.e. for each f_i there exists a finite *separating set* M_{f_i} such that for each $x \in D$ there exists $m \in M_{f_i}$ with $f_i(x) \sqsubseteq m \ll x$, and $\bigsqcup_{i \in I} f_i = \text{id}_D$.

This definition, which differs from the original (Jung 1990) in the use of *strong* finite separation, is nonetheless equivalent to it by Lemma 2 of *op. cit.* Also, as in (Gierz *et al.* 2003), we do not require FS-domains to be pointed. This allows Theorem 5.7 below to be formulated as generally as possible.

Any FS-domain is automatically a continuous dcppo. If D, E are FS-domains then $D \times_s E$ and $D \Rightarrow_s E$ are also FS-domains. Further, if D, E are countably based then so are $D \times_s E$ and $D \Rightarrow_s E$. Thus the categories **FS** and $\omega\mathbf{FS}$ of FS-domains and countably based FS-domains are both cartesian-closed subcategories of **dcppo**. These results are proved in (Jung 1990; Gierz *et al.* 2003).

In contrast to the situation for L-domains, compactly generated function spaces of FS-domains do carry the Scott topology.

Theorem 5.7 *If D, E are FS-domains then $D \Rightarrow_k E = D \Rightarrow_s E = C_{\text{co}}(D, E) = C_{\text{po}}(D, E)$.*

In particular, the inclusions $\mathbf{FS} \hookrightarrow \mathbf{kP}$ and $\omega\mathbf{FS} \hookrightarrow \omega\mathbf{P}$ both preserve the cartesian closed structure.

The first lemma needed in the proof of the theorem is a mild generalization (with identical proof) of Corollary 1.36 of (Jung 1989).

Lemma 5.8 *If D is a dcpo and E a continuous dcpo such that $D \Rightarrow_s E$ is a continuous dcpo, then $f \ll g$ implies $f(x) \ll g(x)$ for all $x \in D$.*

Proof. Suppose $\{e_i\}_{i \in I}$ is a directed family of elements in E with $\bigsqcup_i e_i = g(x)$. We have to show that there exists $i_0 \in I$ such that $e_{i_0} \sqsupseteq f(x)$. Easily, $(\downarrow x)$ and $(\downarrow g(x))$ are continuous retracts of D and E , hence $(\downarrow x) \Rightarrow_s (\downarrow g(x))$ is a continuous retract of $D \Rightarrow_s E$, and therefore a continuous dcpo. For each $h: D \rightarrow E$, let $h': (\downarrow x) \rightarrow (\downarrow g(x))$ denote the image of h under the retraction. Then $f' \ll g'$, for if $\{\psi_j\}_{j \in J}$ is a directed family of functions in $(\downarrow x) \Rightarrow_s (\downarrow g(x))$ with $\bigsqcup_j \psi_j = g'$, then $\{\Psi_j\}_{j \in J}$, defined as

$$\Psi_j(y) = \begin{cases} \psi_j(y) & \text{if } y \sqsubseteq x \\ g(y) & \text{otherwise} \end{cases}$$

is a directed family of functions in $D \Rightarrow_s E$ with $\bigsqcup_j \Psi_j = g$. Thus there exists $j_0 \in J$ such that $\Psi_{j_0} \sqsupseteq f$, and hence $\psi_{j_0} \sqsupseteq f'$.

Now for each $i \in I$, let $c_{e_i}: (\downarrow x) \rightarrow (\downarrow g(x))$ denote the constant function with value e_i . Then $\bigsqcup_i c_{e_i} \sqsupseteq g'$, hence there exists $i_0 \in I$ such that $c_{e_{i_0}} \sqsupseteq f'$, giving

$$e_{i_0} = c_{e_{i_0}}(x) \sqsupseteq f'(x) = f(x),$$

as desired. \square

Lemma 5.9 *If D is an FS-domain and E a continuous dcpo such that $D \Rightarrow_s E$ is a continuous dcpo then $D \Rightarrow_s E$ carries the point open topology.*

Proof. Since D is an FS-domain, there exists a directed set $\{f_i\}_{i \in I}$ of endofunctions each strongly finitely separated from id_D , with finite separating sets M_{f_i} , and $\bigsqcup_i f_i = \text{id}_D$.

Furthermore, since $D \Rightarrow_s E$ is a continuous dcpo, there exists a directed set $\{\psi_j\}_{j \in J}$ of endofunctions such that each $\psi_j \ll h$ and $\bigsqcup_j \psi_j = h$. Then $\{\psi_j \circ f_i\}_{i \in I, j \in J}$ is also a directed set with $\psi_j \circ f_i \ll h$ and $\bigsqcup_j \psi_j \circ f_i = h$. Therefore $\{\hat{\uparrow}(\psi_j \circ f_i)\}_{i \in I, j \in J}$ is a Scott-open neighbourhood basis for $h: D \rightarrow E$. Thus, for each Scott-open neighbourhood \mathcal{U} of h , there exist $i_0 \in I$ and $j_0 \in J$ such that $h \in \uparrow(\psi_{j_0} \circ f_{i_0}) \subseteq \mathcal{U}$. Set $\mathcal{V} = \bigcap_{m \in M_{f_{i_0}}} \langle m, \hat{\uparrow}\psi_{j_0}(m) \rangle$, then $h \in \mathcal{V}$, by Lemma 5.8. We claim that $\mathcal{V} \subseteq \mathcal{U}$. To see this, let $x \in D$ and $h' \in \mathcal{V}$. Then there exists $m \in M_{f_{i_0}}$ with $f_{i_0}(x) \sqsubseteq m \ll x$, and so

$$(\psi_{j_0} \circ f_{i_0})(x) \sqsubseteq \psi_{j_0}(m) \ll h'(m) \sqsubseteq h'(x) .$$

Thus $\mathcal{V} \subseteq \uparrow(\psi_{j_0} \circ f_{i_0}) \subseteq \mathcal{U}$, showing the claim. \square

Proof of Theorem 5.7. If D, E are FS-domains then the conditions of Lemma 5.9 are satisfied, so $D \Rightarrow_s E$ carries the point open topology which, by Lemma 5.5, coincides with the compact open topology. So $C_{\text{co}}(D, E)$ carries the Scott topology, and is thus compactly generated. Whence $D \Rightarrow_k E = C_{\text{co}}(D, E) = D \Rightarrow_s E$. \square

Theorem 5.7 requires both domain and codomain of the function space to be FS-domains. In contrast, Proposition 3.7 asserts that the compactly generated exponential $X \Rightarrow_k \mathbb{S}$ carries the Scott topology for every compactly generated space X . We end the paper by considering to what extent this property generalises to continuous dcpos other than Sierpinski space \mathbb{S} . Clearly it does not always hold since, by Proposition 5.3, the property fails when \mathbb{S} is replaced by the five element pointed poset L_2 of Fig. 1.

Theorem 5.10 *If X is compactly generated and E is a continuous dcpo with binary infima then $X \Rightarrow_k E = X \Rightarrow_s E$.*

For the proof, we need a lemma which is part of the domain-theoretic folklore.

Lemma 5.11 *If a continuous dcpo has binary infima then it has infima for all nonempty compact subsets.*

As we could only find an indirect proof in the literature, (Schalk 1993, Lemma 7.14), it seems worth giving a direct argument.

Proof. Suppose D has binary, and therefore nonempty finite meets. Let $K \subseteq D$ be nonempty and compact. Then $\downarrow K = \{x \in D \mid \forall y \in K. x \ll y\}$ is nonempty, as $K \subseteq \bigcup_{x \in D} \hat{\uparrow}x$, so there exists a nonempty finite $F \subseteq D$ such that $K \subseteq \bigcup_{x \in F} \hat{\uparrow}x$, hence $\bigwedge F \ll K$. We claim that $\downarrow K$ is directed.

To see this let $a, b \in \downarrow K$. Then for each $x \in K$, there exists $c_x \in D$ such that $a, b \sqsubseteq c_x \ll x$. Thus $K \subseteq \bigcup_{x \in K} \hat{\uparrow}c_x$, and so there exists a non empty finite $F \subseteq K$ such that $K \subseteq \bigcup_{x \in F} \hat{\uparrow}c_x$. But then $\bigwedge_{x \in F} c_x$ fulfills $a, b \sqsubseteq \bigwedge_{x \in F} c_x \ll K$, showing the claim.

Now let $x \sqsubseteq K$. Then $x = \bigsqcup \downarrow x = \bigsqcup (\downarrow x \cap \downarrow K)$. Thus $x \sqsubseteq \bigsqcup \downarrow K$, and so $\bigsqcup \downarrow K$ is the greatest lower bound of K , as required. \square

Proof of Theorem 5.10. By Theorem 3.11, $X \Rightarrow_k E$ is a monotone convergence space. Thus we just need to verify that every Scott-open subset of $X \Rightarrow_k E$ is indeed open.

Let $W \subseteq X \Rightarrow_k E$ be Scott-open. Consider any continuous $p: K \rightarrow (X \Rightarrow_k E)$ with K compact Hausdorff. We must show that $p^{-1}(W) \subseteq K$ is open. Suppose we have $k \in K$ with $p(k) \in W$. We must find a neighbourhood of k contained in $p^{-1}(W)$.

Let D be the set of compact neighbourhoods of k , ordered by reverse inclusion. K is compact Hausdorff, hence locally compact, so D is directed. For every $L \in D$, define $h_L: X \rightarrow E$, by

$$h_L(x) = \prod_{z \in L} p(z)(x).$$

Then h_L is continuous because it arises as a composite of continuous functions

$$X \xrightarrow{\tilde{p}} C_{\text{co}}(K, E) \longrightarrow C_{\text{co}}(L, E) \xrightarrow{M_L} E,$$

the components of which we now describe. The map \tilde{p} is obtained by the following manipulations.

$$\begin{array}{lll} p: K & \longrightarrow & (X \Rightarrow_k E) \\ K \times_k X & \longrightarrow & E & \text{exponential transpose in } \mathbf{kTop} \\ K \times X & \longrightarrow & E & \text{by Proposition 3.3, using } K \text{ core compact} \\ \tilde{p}: X & \longrightarrow & C_{\text{co}}(K, E) & \text{exponential transpose in } \mathbf{Top}. \end{array}$$

The map $C_{\text{co}}(K, E) \rightarrow C_{\text{co}}(L, E)$ is induced by the inclusion $L \subseteq K$. Finally M_L is the function $M_L(f) = \prod_{z \in L} f(z)$. This is continuous because if $y \ll M_L(f)$ then there exists y' with $y \ll y' \ll M_L(f)$. Hence $\langle L, \uparrow y' \rangle$ is a neighbourhood of f in $C_{\text{co}}(L, E)$ satisfying $y \ll M_L(g)$, for all $g \in \langle L, \uparrow y' \rangle$.

Easily $L' \supseteq L \in D$ implies $h_{L'} \sqsubseteq h_L \in X \Rightarrow_s E$, so $H = \{h_L \mid L \in D\}$ is a directed subset of $X \Rightarrow_s E$. We show that $\bigsqcup H = p(k)$.

For every $L \in D$, we have $k \in L$, so $h_L(x) = \prod_{z \in L} p(z)(x) \sqsubseteq p(k)(x)$. Therefore $\bigsqcup H \sqsubseteq p(k)$.

Conversely, we show that $p(k) \sqsubseteq \bigsqcup H$, i.e. $p(k)(x) \sqsubseteq (\bigsqcup H)(x)$, for all $x \in X$. Take any $x \in X$ and $y \ll p(k)(x)$. By the continuity of p , the set $U = \{z \mid y \ll p(z)(x)\} \subseteq K$ is open. Also $k \in U$. By local compactness, there exists a compact neighbourhood $L \ni k$ with $L \subseteq U$. Then $h_L(x) = \prod_{z \in L} p(z)(x) \supseteq y$. Thus indeed $y \sqsubseteq \bigsqcup (H(x)) = (\bigsqcup H)(x)$.

In summary, we have directed $H \subseteq X \Rightarrow_k E$ with $\bigsqcup H = p(k) \in W$. As W is Scott-open, there exists $L \in D$ such that $h_L \in W$. Since L is a neighbourhood of k , we just need to show that $L \subseteq p^{-1}W$. So consider any $z \in L$. Then, we have $p(z) \supseteq h_L \in W$. Thus indeed $p(z) \in W$, since W is upper-closed. \square

We remark that a special case of Theorem 5.10 follows more easily from existing results in the domain-theoretic literature. It is known that if X is a locally compact topological space and E is a bounded-complete continuous dcppo then $C_{\text{co}}(X, E)$ carries the Scott topology and is itself a bounded-complete continuous dcppo, see Proposition II-4.6 of (Gierz *et al.* 2003). So, in this case, the coincidence of the spaces $X \Rightarrow_k E$ and $X \Rightarrow_s E$ follows as a consequence of Propositions 3.2 and 3.5. We do not know if, more generally, $C_{\text{co}}(X, E)$ carries the Scott topology, for locally compact X , when E merely has binary infima. It certainly does not hold in general that $X \Rightarrow_s E$ is a continuous




Fig. 2.

dcpo in this case. A counterexample is the space $U_\omega \Rightarrow_s U_\omega$, where U_ω is the well-known non-bifinite ω -algebraic dcppo from Fig. 2. Although U_ω has binary infima, the dcppo $U_\omega \Rightarrow_s U_\omega$ is not continuous. Thus, from a traditional domain-theoretic viewpoint, the Scott topology on the function space $U_\omega \Rightarrow_s U_\omega$ is poorly behaved. In contrast, from a compactly generated viewpoint, the Scott topology is well behaved in this case. By Theorem 5.10, the function spaces $U_\omega \Rightarrow_k U_\omega$ and $U_\omega \Rightarrow_s U_\omega$ coincide. Moreover, by Proposition 3.6 (and Lemma 5.5), they have the compact open (equivalently point open) topology, and this is countably based.

6. Discussion

In this paper we have demonstrated that the topological (pre)domains of (Simpson 2003) fit naturally into the world of convenient topology. As argued in *op. cit.*, topological predomains overcome the limitations, discussed in Section 1, of traditional domain theory. Moreover, the larger collection of compactly generated (pre)domains investigated in the present paper provides a natural topological generalization of the dcpo-based world of traditional domain theory.

It is appropriate to question the use of compactly generated spaces in this paper. Even in algebraic topology, it is hard to give *a priori* justification for taking the notion of compactly generated space as basic. From a domain-theoretic perspective, the motivation is even less clear. In particular, the choice of compact Hausdorff spaces as the generating spaces in Definition 3.1 seems utterly arbitrary.

In fact, compactly generated spaces form just one of many cartesian closed subcategories of **Top**. Arguably, a more natural subcategory is the category of *core compactly*

generated spaces introduced in (Day 1972), which is the largest cartesian closed subcategory of **Top** obtainable using the general theory of (Day 1972; Escardó *et al.* 2004). This category properly includes the category of compactly generated spaces (Isbell 1987). It seems likely that the results of the present paper should generalise to taking core compactly generated monotone convergence space as a notion of predomain, and other variants should be possible too.

Alternatively, some might prefer to carry out an analogous generalization of domain theory within a cartesian closed supercategory of **Top**, such as Scott’s category of equilogical spaces (Bauer *et al.* 2004), or one of the categories of “convergence” or “filter” spaces, see e.g. (Wyler 1974; Hyland 1979). In fact, to some extent, Reinhold Heckmann has already embarked upon such a programme. In (Heckmann 2003), he develops a convergence space variant of the notion of monotone convergence space, and establishes results analogous to our Propositions 3.14, 5.1 and 5.2 for that notion.

It is a pleasing fact that apparent differences between the subcategory and supercategory approaches disappear if attention is restricted to qcb spaces. As shown in (Menni and Simpson 2002; Escardó *et al.* 2004), the category **QCB** lives, via structure-preserving embeddings, in all the principal cartesian closed subcategories of **Top**, and also in Scott’s category of equilogical spaces. (It is the latter embedding that forms the basis of the connections with realizability semantics mentioned in Section 1.) Analogous embeddings of **QCB** in categories of convergence spaces have not been established, but are expected.

Seemingly, **QCB** is an inevitable category, occurring within any sufficiently general approach to combining cartesian closedness and topology. In the authors’ view, it is the category of paramount interest for the semantics of computation. For example, the size restriction naturally expresses the requirement that data should be representable by a sequence of discrete approximations (Weihrauch 2000; Schröder 2003).

Each of the larger categories embedding **QCB** offers its own valuable perspective on qcb spaces. In particular, as demonstrated in this paper, the approach via compactly generated spaces provides a good framework for relating topological (pre)domains and traditional domain theory. Thus, even though the notion of compactly generated space apparently lacks intrinsic motivation, such spaces do nonetheless provide a useful bridge between traditional topological domain theory and its topological generalizations.

Acknowledgements

We have benefited from discussions with Martín Escardó, Jimmie Lawson and Thomas Streicher. We also thank the editors of this volume for their patience and encouragement.

References

- S. Abramsky and A. Jung. Domain theory. In *Handbook of Logic in Computer Science, Vol. 3*, pages 1–168. OUP, 1994.
- I. Battenfeld. *A Category of Topological Predomains*. Diploma Thesis, TU Darmstadt, 2004.
- A. Bauer, L. Birkedal, and D.S. Scott. Equilogical spaces. *Theoretical Computer Science*, 315:35–59, 2004.

- R. Brown. Ten topologies for $X \times Y$. *Quart. J. Math. Oxford Ser.*, 14:303–319, 1963.
- R. Brown. Function spaces and product topologies. *Quart. J. Math. Oxford Ser.*, 15:238–250, 1964.
- B.J. Day. A reflection theorem for closed categories. *J. Pure Appl. Algebra*, 2:1–11, 1972.
- M.H. Escardó, J.D. Lawson, and A.K. Simpson. Comparing cartesian closed categories of core compactly generated spaces. *Topology and its Applications*, 143:105–145, 2004.
- M.H. Escardó. Notes on compactness. Preprint, 2005.
- G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M.W. Mislove, and D.S. Scott. *Continuous Lattices and Domains*. Number 93 in Encyclopedia of Mathematics and its Applications. CUP, 2003.
- R. Heckmann. A non-topological view of dcpos as convergence spaces. *Theoretical Computer Science*, 305:159–186, 2003.
- J.M.E. Hyland. Filter spaces and continuous functionals. *Annals of Mathematical Logic*, 16:101–143, 1979.
- J. Isbell. A distinguishing example in k -spaces. *Proc. Amer. Math. Soc.*, 100:593–594, 1987.
- A. Jung. *Cartesian closed categories of domains*, volume 66 of *CWI Tracts*. CWI, Amsterdam, 1989.
- A. Jung. The classification of continuous domains. In *Proceedings 5th IEEE Symposium on Logic in Computer Science*, pages 35–40, 1990.
- P.Th. Lambrinos and B. Papadopoulos. The (strong) Isbell topology and (weakly) continuous lattices. In *Continuous lattices and their applications (Bremen, 1982)*, pages 191–211, Lecture Notes in Pure and Appl. Math., 101, Dekker, New York, 1985.
- J.R. Longley and A.K. Simpson. A uniform account of domain theory in realizability models. *Math. Struct. in Comp. Sci.*, 7:469–505, 1997.
- E. Michael. \aleph_0 -spaces. *J. Math. Mech.*, 15:983–1002, 1966.
- M. Menni and A.K. Simpson. Topological and limit space subcategories of countably-based equilogical spaces. *Math. Struct. in Comp. Sci.*, 12, 2002.
- W.K.-S. Phoa. Effective domains and intrinsic structure. In *Proc. 5th Annual Symposium on Logic in Computer Science*, pages 366–377, 1990.
- G. Preuss. *Foundations of topology: an approach to convenient topology*. Kluwer, 2002.
- B. Reus and Th. Streicher. General synthetic domain theory — a logical approach. *Math. Struct. in Comp. Sci.*, 9:177–223, 1999.
- A. Schalk. *Algebras for Generalized Power Constructions*. PhD thesis, Technischen Hochschule Darmstadt, 2003.
- M. Schröder. *Admissible Representations for Continuous Computations*. PhD thesis, FernUniversität, Hagen, 2003.
- D.S. Scott. Data types as lattices. *SIAM J. Comput.*, 5:522–587, 1976.
- A.K. Simpson. Towards a convenient category of topological domains. In *Proceedings of thirteenth ALGI Workshop*. RIMS, Kyoto University, 2003.
- N.E. Steenrod. A convenient category of topological spaces. *Michigan Math. J.*, 14:133–152, 1967.
- K. Weihrauch. *Computable Analysis*. Springer, 2000.
- O. Wyler. Filter space monads, regularity, completions. In *Proceedings TOPO 72 — General Topology and its Applications*, pages 591–637. Springer LNM 378, 1974.
- O. Wyler. Dedekind complete posets and scott topologies. In *Continuous Lattices, Proceedings of the Conference on Topological and Categorical Aspects of Continuous Lattices, Bremen 1979*, pages 384–389. Springer LNM 871, 1981.