

Noncommutativity as a colimit

Benno van den Berg* and Chris Heunen†

July 20, 2010

Abstract

Every partial algebra is the colimit of its total subalgebras. We prove this result for partial Boolean algebras (including orthomodular lattices) and the new notion of partial C*-algebras (including noncommutative C*-algebras), and variations such as partial complete Boolean algebras and partial AW*-algebras. The first two results are related by taking projections. As corollaries we find extensions of Stone duality and Gelfand duality. Finally, we investigate the extent to which the Bohrification construction [9], that works on partial C*-algebras, is functorial.

1 Introduction

This paper is intended as a contribution to the Bohrification programme [9], which tries to give a mathematically precise expression to Bohr's doctrine of classical concepts, saying that a quantum mechanical system is to be understood through its classical fragments. On Bohr's view, as understood within this programme, quantum mechanical systems do not allow a 'global' interpretation as a classical system, but they do so 'locally'. The programme essentially makes two claims about these classical 'snapshots'. Firstly, it is only through these snapshots that the behaviour of a system and physical reality can be understood. Secondly, these snapshots contain all the information about the system that is physically relevant.

The main result of [9] (inspired by earlier work of Butterfield, Isham and Döring [5]) is that the collection of these classical snapshots can be seen as forming a single classical system in a suitable topos – its so-called Bohrification; we will briefly recall the details of this construction in Section 7. The implication of this result is that a quantum mechanical system can be seen as a classical one, if one agrees that nothing physically relevant is lost by considering classical snapshots and if one is willing to change the logic from a classical into an intuitionistic one.

*Technische Universität Darmstadt, Fachbereich Mathematik.

†Oxford University Computing Laboratory, supported by the Netherlands Organisation for Scientific Research (NWO). Part of this work was performed while the author visited the IQI at Caltech.

Our conceptual contributions to the Bohrification programme are twofold. First, we contend that the programme is most naturally seen as concerned with *partial algebras*, *i.e.* sets on which algebraic structure is only partially defined. Such partial algebras are equipped with a binary (‘commeasurability’) relation, which holds between two elements whenever they can be elements of a single classical snapshot. In this setting, commeasurability is often also called commutativity, since in typical examples commeasurability means commutativity in a totally defined algebraic structure. We will show that a meaningful theory of such partial algebras can be developed. In particular, and this is our second contribution, we will indicate why Bohrification makes sense for partial algebras and use this to investigate the functorial aspects of this construction.

The idea to consider partial algebras is not new. In fact, in their classic paper [14], Kochen and Specker use the language of partial algebras to state their famous result, saying that many algebras occurring in quantum mechanics cannot be embedded (as partial algebras) into commutative ones (see also [17]). Their interpretation of this fact as excluding a hidden variable interpretation of quantum mechanics has remained somewhat controversial. In the Bohrification programme, this result is taken just as a mathematical confirmation of the view that a quantum mechanical systems do not allow for global interpretations as classical systems.

The result by Kochen and Specker, together with the fact that Bohrification works for partial algebras, indicates that the Bohrification programme is most naturally developed in the context of partial algebras. This led us to develop a new notion of ‘partial C*-algebra’, to the study of which most of this paper is devoted. Our main technical result is that such a partial C*-algebra is the colimit of its total (commutative) subalgebras, which explains the relation between a partial algebra and its classical snapshots in categorical terms.

In more detail, the contents of this paper are as follows. First we consider Kochen and Specker’s notion of a partial Boolean algebra, as a kind of toy example, and prove our main result for them in Section 2, as well as for the variation of partial complete Boolean algebras. This also makes precise the widespread intuition that an orthomodular lattice is an amalgamation of its Boolean blocks. In this simple setting the theorem already has interesting corollaries, as it allows us to derive an adjunction extending Stone duality in Section 3. But our main interest lies with partial C*-algebras, which Section 4 studies; we also prove the main result for variations, such as AW*-algebras and Rickart C*-algebras in that section. This enables an adjunction extending Gelfand duality and puts the Kochen-Specker theorem in another light in Section 5. The two parallel settings of Boolean algebras and C*-algebras are related by taking projections, as Section 6 discusses. Finally, Section 7 shows that the Bohrification works for partial C*-algebras as well: we investigate its functorial properties, and conclude that its essence is in fact (a two-dimensional version of) the colimit theorem.¹

¹Both the categories of Boolean algebras and of commutative C*-algebras are algebraic, *i.e.* monadic over the category of sets [10]. We expect that the definitions and results of the present article can be extended to a more general theory of partial algebra, but refrain from doing so because the two categories mentioned are our main motivation.

2 Partial Boolean algebras

We start with recalling the definition of a partial Boolean algebra, as introduced by Kochen en Specker [14].

Definition 1 A *partial Boolean algebra* consists of a set A with

- a reflexive and symmetric binary (*commensurability*) relation $\odot \subseteq A \times A$;
- elements $0, 1 \in A$;
- a (total) unary operation $\neg: A \rightarrow A$;
- (partial) binary operations $\wedge, \vee: \odot \rightarrow A$;

such that every set $S \subseteq A$ of pairwise commensurable elements is contained in a set $T \subseteq A$, whose elements are also pairwise commensurable, and on which the above operations determine a Boolean algebra structure.²

A morphism of partial Boolean algebras is a function that preserves commensurability and all the algebraic structure, whenever defined. We write **PBoolean** for the resulting category.

Clearly, a partial Boolean algebra whose commensurability relation is total is nothing but a Boolean algebra. For another example, if we declare two elements a, b of an orthomodular lattice to be commensurable when $a = (a \wedge b) \vee (a \wedge b^\perp)$, as is standard, any orthomodular lattice is seen to be a partial Boolean algebra. In this case the above observation about totality becomes a known fact: an orthomodular lattice is Boolean if and only if it is commutative [13].

We introduce some notation and terminology. If A is a partial Boolean algebra and S is subset of pairwise commensurable elements, then there must be a *smallest* set T of pairwise commensurable elements that contains S and which has a Boolean algebra structure determined by the operations of A : it has to consist of the values of Boolean expressions built from elements from S . We denote it by $A\langle S \rangle$.

Given a partial Boolean algebra A , the collection of its commensurable subalgebras $\mathcal{C}(A)$ is partially ordered by inclusion. In fact, \mathcal{C} is a functor **PBoolean** \rightarrow **POrder** to the category of posets and monotone functions. Regarding posets as categories, $\mathcal{C}(A)$ gives a diagram in the category **PBoolean**. The following proposition lists some easy properties of this diagram.

Proposition 2 *Let A be a partial Boolean algebra.*

- (a) *The least element of the poset $\mathcal{C}(A)$ is $A\langle 0 \rangle = A\langle 1 \rangle = \{0, 1\}$.*
- (b) *The poset $\mathcal{C}(A)$ is filtered if and only if A itself is Boolean. In that case, A is the largest element of the poset $\mathcal{C}(A)$.*
- (c) *The atoms of $\mathcal{C}(A)$ are $A\langle a \rangle = \{0, a, \neg a, 1\}$ for nontrivial $a \in A$. □*

²Note that this means that T must contain 0 and 1 and has to be closed under \neg, \wedge and \vee .

Remark 3 One can show that $\mathcal{C}(A)$ is a directed complete partial order (dcpo), which is algebraic, and is such that for every compact element x the downset $\downarrow x$ is dually isomorphic to a finite partition lattice. The main result of the paper [6] suggests that every such dcpo is the $\mathcal{C}(A)$ of a unique partial Boolean algebra A . Whether similar results hold for partial C^* -algebras remains to be seen.

We are now ready to prove the first version of our main result.

Theorem 4 *Every partial Boolean algebra is a colimit of its (finitely generated) Boolean subalgebras.*


PROOF Let A be a partial Boolean algebra, and consider its diagram of (finitely generated) commensurable subalgebras C . Define functions $i_C: C \rightarrow A$ by the inclusions; these are morphisms of **PBoolean** by construction. Moreover, they form a cocone; we will prove that this cocone is universal. If $f_C: C \rightarrow B$ is another cocone, define a function $m: A \rightarrow B$ by $m(a) = f_{A\langle a \rangle}(a)$. It now follows from the assumption that the f_C are morphisms of **PBoolean** that m is a well-defined morphism, too. For example, if $a \odot b$, then also $m(a) \odot m(b)$, since $A\langle a, b \rangle$ is a Boolean subalgebra and

$$\begin{aligned} m(a) &= f_{A\langle a \rangle}(a) = f_{A\langle a, b \rangle}(a), \\ m(b) &= f_{A\langle b \rangle}(b) = f_{A\langle a, b \rangle}(b), \end{aligned}$$

because the f_C form a cocone. One easily verifies that $f_C = m \circ i_C$, and that m is the unique such morphism. \square

Kalmbach’s “Bundle Lemma” [13] gives sufficient conditions for a family of Boolean algebras to combine into a partial Boolean algebra, so that it could be regarded as a converse of the previous theorem.

Notice that the morphisms for **PBoolean** are the weakest ones for which the previous theorem holds. For example, the mediating morphism m in the proof of the previous theorem need not be a homomorphism of orthomodular lattices. For a counterexample, consider the function



given by $m(0) = 0, m(a) = m(b) = c, m(a^\perp) = m(b^\perp) = c^\perp, m(1) = 1$. It preserves $0, 1, \perp$ and \leq . The only Boolean subalgebras of the domain A are $A\langle 0 \rangle, A\langle a \rangle$ and $A\langle b \rangle$, and m preserves \wedge when restricted to those. However, $m(a \wedge b) = m(0) = 0 \neq c = c \wedge c = m(a) \wedge m(b)$.

It follows from the previous theorem that the partial Boolean algebra with a prescribed poset of Boolean subalgebras is unique up to isomorphism (of partial Boolean algebras). To actually reconstruct a partial Boolean algebra from its

Boolean subalgebras, this should be complemented by a description of colimits in the category **PBoolean**, as we now discuss.

The coproduct of a family A_i of partial Boolean algebras is got by taking their disjoint union, identifying all the elements 0_i , and identifying all the elements 1_i . Notice that elements from different summands A_i are never commensurable in the coproduct. In particular, the initial object $\mathbf{0}$ is the partial Boolean algebra $\{0, 1\}$ with two distinct elements.

Incidentally, **PBoolean** is complete. Products and equalizers of partial Boolean algebras are constructed as in the category of sets; products have commensurability and algebraic structure defined componentwise, and equalisers have subalgebra structure. Hence the limit of a diagram of Boolean algebras is the same in the categories of Boolean algebras and of partial Boolean algebras. In particular, the terminal object $\mathbf{1}$ is the partial Boolean algebra with a single element $0 = 1$.

Coequalizers are harder to describe constructively, but the following theorem proves they do exist.

Theorem 5 *The category **PBoolean** is complete and cocomplete.*

PROOF We are to show that **PBoolean** has coequalizers, *i.e.* that the diagonal functor $\Delta: \mathbf{PBoolean} \rightarrow \mathbf{PBoolean}^{(\cdot \rightrightarrows \cdot)}$ has a left adjoint. Since we already know that **PBoolean** is complete and Δ preserves limits, Freyd's adjoint functor theorem shows that it suffices if the following solution set condition is satisfied [15, V.6]. For each $f, g: A \rightarrow B$ in **PBoolean** there is a set-indexed family $h_i: B \rightarrow Q_i$ such that $h_i f = h_i g$, and if $h f = h g$ then h factorizes through some h_i .

Take the collection of $h_i: B \rightarrow Q_i$ to comprise all 'quotients', *i.e.* (isomorphism classes of) surjections h_i of partial Boolean algebras such that $h_i f = h_i g$. This collection is in fact a set. The proof is finished by observing that every morphism $h: B \rightarrow Q$ of partial Boolean algebras factors through (a surjection onto) its set-theoretical image, which is a partial Boolean subalgebra of Q , inheriting commensurability from B and algebraic operations from Q . \square

2.1 Variations

Results similar to those above hold for many classes of Boolean algebras, such as complete or countably complete Boolean algebras. For example, the former variation can be defined as follows.

Definition 6 A *partial complete Boolean algebra* consists of a partial Boolean algebra together a (partial) operation

$$\bigvee: \{X \subseteq A \mid X \times X \subseteq \odot\} \rightarrow A$$

such that every set $S \subseteq A$ of pairwise commensurable elements is contained in a set $T \subseteq A$, whose elements are also pairwise commensurable, and on which the above operations determine a complete Boolean algebra structure.³

³So T is not only closed under \neg, \wedge and \vee , but also under \bigvee .

A morphism of partial complete Boolean algebras is a function that preserves commensurability and all the algebraic structure, including \bigvee , whenever defined. We write **PCBoolean** for the resulting category.

A version of our main theorem also holds for such partial complete Boolean algebras.

Theorem 7 *Every partial complete Boolean algebra is a colimit of its complete Boolean subalgebras.*

PROOF Completely analogous to Theorem 4. □

3 Stone duality

The full subcategory of **PBoolean** consisting of (total) Boolean algebras is just the category **Boolean** of Boolean algebras and their homomorphisms. This category is dual to the category of Stone spaces and continuous functions via Stone duality [10]:

$$\mathbf{Boolean} \begin{array}{c} \xrightarrow{\Sigma} \\ \xleftarrow{\sim} \\ \xleftarrow{\mathbf{Loc}(-, \{0,1\})} \end{array} \mathbf{Stone}^{\text{op}}, \quad (1)$$

where $\Sigma(A)$ is the Stone spectrum of a Boolean algebra A . The dualizing object $\{0,1\}$ is both a locale and a (partial) Boolean algebra; recall that it is in fact the initial partial Boolean algebra $\mathbf{0}$.

One might expect that the category of partial Boolean algebras enters Stone duality (1), and indeed the colimit theorem, Theorem 4, enables us to prove the following extension.

Proposition 8 *There is a reflection*

$$\mathbf{PBoolean} \begin{array}{c} \xrightarrow{K} \\ \xleftarrow{\perp} \\ \xleftarrow{\mathbf{Loc}(-, \{0,1\})} \end{array} \mathbf{Stone}^{\text{op}},$$

in which the functor K is determined by $K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C)$.

PROOF Let A be a partial Boolean algebra and X a Stone space. Then there are bijective correspondences:

$$\begin{array}{l} \underline{\underline{f: K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C) \rightarrow X}} \quad (\text{in } \mathbf{Stone}^{\text{op}}) \\ \underline{\underline{\forall_{C \in \mathcal{C}(A)} \cdot f_C: \Sigma(C) \rightarrow X}} \quad (\text{in } \mathbf{Stone}^{\text{op}}) \\ \underline{\underline{\forall_{C \in \mathcal{C}(A)} \cdot g_C: C \rightarrow \mathbf{Loc}(X, \{0,1\})}} \quad (\text{in } \mathbf{Boolean}) \\ \underline{\underline{g: A \rightarrow \mathbf{Loc}(X, \{0,1\})}} \quad (\text{in } \mathbf{PBoolean}). \end{array}$$

The first correspondence holds by definition of limit, the middle correspondence holds by Stone duality (1), and the last correspondence holds by Theorem 4.

Since all correspondences are natural in A and X , this establishes the adjunction $K \dashv \mathbf{Loc}(-, \{0, 1\})$. Finally, since a Boolean algebra is trivially a colimit of itself in $\mathbf{PBoolean}$, the adjunction is a reflection. \square

Theorem 9 *The reflection $K \dashv \mathbf{Loc}(-, \{0, 1\})$ extends Stone duality, i.e. the following diagram commutes (serially).*

$$\begin{array}{ccc}
 & \mathbf{Boolean} & \\
 & \swarrow & \searrow \\
 \mathbf{PBoolean} & & \mathbf{Stone}^{\text{op}} \\
 & \swarrow & \searrow \\
 & \mathbf{Stone}^{\text{op}} & \\
 & \swarrow & \searrow \\
 \mathbf{PBoolean} & & \mathbf{Stone}^{\text{op}}
 \end{array}$$

$\mathbf{Loc}(-, \{0, 1\})$ (top left arrow), Σ (top right arrow), \sim (middle arrow), K (middle arrow), \perp (bottom arrow), $\mathbf{Loc}(-, \{0, 1\})$ (bottom arrow)

PROOF If A is a Boolean algebra, it is the initial element in the diagram $\mathcal{C}(A)^{\text{op}}$ by Proposition 2(b). Hence $K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C) = \Sigma(A)$. \square

Corollary 10 *Boolean algebras form a reflective full subcategory of the category of partial Boolean algebras, i.e. the inclusion $\mathbf{Boolean} \hookrightarrow \mathbf{PBoolean}$ has a left adjoint $L: \mathbf{PBoolean} \rightarrow \mathbf{Boolean}$.*

PROOF The adjunctions of the previous theorem compose, giving the required left adjoint as $L = \mathbf{Loc}(-, \{0, 1\}) \circ K$. \square

4 Partial C*-algebras

The definitions of partial C*-algebras and their morphisms closely resemble those of partial Boolean algebras. Indeed, both are instances of the partial algebras of Kochen and Specker, over the fields \mathbb{Z}_2 and \mathbb{C} , respectively. However, partial C*-algebras also have to account for a norm and involution, calling for some changes that we now spell out.

Definition 11 A *partial C*-algebra* consists of a set A with

- a reflexive and symmetric binary (*commensurability*) relation $\odot \subseteq A \times A$;
- elements $0, 1 \in A$;
- a (total) involution $*$: $A \rightarrow A$;
- a (total) function \cdot : $\mathbb{C} \times A \rightarrow A$;
- a (total) function $\|-\|$: $A \rightarrow \mathbb{R}$;
- (partial) binary operations $+, \cdot$: $\odot \rightarrow A$;

such that every set $S \subseteq A$ of pairwise commensurable elements is contained in a set $T \subseteq A$, whose elements are also pairwise commensurable, and on which the above operations determine the structure of a commutative C*-algebra.⁴

⁴This entails that T contains 0 and 1, is closed under all algebraic operations, and is norm-complete.

It follows from the last condition in the definition of a partial C^* -algebra that commensurable elements in a partial C^* -algebra have to commute. In fact, as with partial Boolean algebras, partial C^* -algebras whose commensurability relation are total are nothing but *commutative* C^* -algebras.

Also, if A is a partial C^* -algebra and S is subset of pairwise commensurable elements, then there must be a *smallest* set T of pairwise commensurable elements that contains S and which has a commutative C^* -algebra structure determined by the operations of A : simply take the intersection of all such sets T . Alternatively, one can construct it as the set of those elements in A that are limits of sequences whose terms are algebraic expressions involving the elements of S . We denote it by $A\langle S \rangle$.

The reader may be tempted to believe that every noncommutative C^* -algebra can be regarded as a partial C^* -algebra by declaring that $a \odot b$ holds whenever a and b commute, but that would be incorrect. The reason for this is that we have required \odot to be reflexive, so that $aa^* = a^*a$ holds for every element a in a partial C^* -algebra. Now, an element a such that $aa^* = a^*a$ holds is called *normal* and it is not the case that every element in a C^* -algebra is normal.

What is true, however, is that one may regard the collection of normal elements of a C^* -algebra as a partial C^* -algebra by declaring two elements to be commensurable whenever they commute. In fact, taking normal elements is part of a functor, if we consider the following class of morphisms of partial C^* -algebras.

Definition 12 A *partial $*$ -morphism* is a (total) function $f: A \rightarrow B$ between partial C^* -algebras such that:

- $f(a) \odot f(b)$ for commensurable $a, b \in A$;
- $f(ab) = f(a)f(b)$ for commensurable $a, b \in A$;
- $f(a + b) = f(a) + f(b)$ for commensurable $a, b \in A$;
- $f(a + ib) = f(a) + if(b)$ for self-adjoint $a, b \in A$;
- $f(za) = zf(a)$ for $z \in \mathbb{C}$ and $a \in A$;
- $f(a)^* = f(a^*)$ for $a \in A$;
- $f(1) = 1$.

Partial C^* -algebras and partial $*$ -morphisms organize themselves into a category denoted by **PCstar**.⁵

Before we embark on proving that taking normal parts provides a functor from the category of C^* -algebras to the category of partial C^* -algebras, recall that an element a of a C^* -algebra is called self-adjoint when $a = a^*$, and that any element can be written uniquely as a linear combination $a = a_1 + ia_2$ of two self-adjoint elements $a_1 = \frac{1}{2}(a + a^*)$ and $a_2 = \frac{1}{2i}(a - a^*)$.

⁵Most results hold for nonunital C^* -algebras, but for convenience we consider unital ones.

Proposition 13 *There is a functor $N: \mathbf{Cstar} \rightarrow \mathbf{PCstar}$ which sends every C^* -algebra to its normal part*

$$N(A) = \{a \in A \mid aa^* = a^*a\},$$

which can be considered as a partial C^ -algebra by saying that $a \odot b$ holds whenever a and b commute. Moreover, N is faithful and reflects isomorphisms and identities.*

PROOF The action of N is well-defined on objects, since a subalgebra of a C^* -algebra generated by a set S is commutative iff the elements of S are normal and commute pairwise.

On morphisms, N acts by restriction and corestriction. To see that it has the properties the proposition claims it has, one uses the identity $a = a_1 + ia_2$. For example, suppose $N(f)$ is surjective for a $*$ -morphism $f: A \rightarrow B$ and let $b \in B$. Then there are $a_1, a_2 \in N(A)$ with $f(a_i) = b_i$. Hence $f(a_1 + ia_2) = b$, so that f is surjective. Similarly, if $N(f)$ is injective, suppose that $f(a) = f(a')$. Then $f(a_i) = f(a'_i)$, and hence $a_i = a'_i$, so that $a = a'$ and f is injective. Now, isomorphisms in $(\mathbf{P})\mathbf{Cstar}$ are bijective (partial) $*$ -morphisms. Hence f is an isomorphism when $N(f)$ is. \square

So therefore one way of thinking about a partial C^* -algebra is as axiomatizing the normal part of a C^* -algebra. Of course, we could have decided to drop the requirement that \odot is reflexive, so that every C^* -algebra would also be a partial C^* -algebra, with commensurability given by commutativity. We haven't done this for various reasons. First of all, we would like to have a notion given in terms of the physically relevant data. On Bohr's philosophy, which we adopt in this paper, the physically relevant information is contained in the normal part of a C^* -algebra. Related to this is the fact that the Bohrification functor (which we will study in Section 7) only takes the normal part of a C^* -algebra into account. Secondly, we wish to have a result saying how a partial C^* -algebra is determined by its commutative subalgebras analogous to our result for partial Boolean algebras. With our present definition we do indeed have this result (it is Theorem 15 below), as we will now explain.

Denote by $\mathcal{C}: \mathbf{PCstar} \rightarrow \mathbf{POrder}$ the functor assigning to a partial C^* -algebra A the collection of its commensurable (*i.e.* commutative total) subalgebras $\mathcal{C}(A)$, partially ordered by inclusion. One immediately derives similar properties for the diagram $\mathcal{C}(A)$ in the category \mathbf{PCstar} as for partial Boolean algebras.

Proposition 14 *Let A be a partial C^* -algebra.*

(a) *The least element of the poset $\mathcal{C}(A)$ is $A\langle 0 \rangle = A\langle 1 \rangle = \{z \cdot 1 \mid z \in \mathbb{C}\}$.*

(b) *The poset $\mathcal{C}(A)$ is filtered if and only if A is a commutative C^* -algebra. In that case, A is the largest element of the poset $\mathcal{C}(A)$. \square*

We now prove the C^* -algebra version of our main result.

Theorem 15 *Every partial C*-algebra is a colimit of its (finitely generated) commutative C*-subalgebras.*

PROOF Let A be a partial C*-algebra, and consider its diagram $\mathcal{C}(A)$ of (finitely generated) commutative C*-subalgebras C . Defining functions $i_C: C \rightarrow A$ by the inclusions yields a cocone in **PCstar**; we will prove that this cocone is universal. If $f_C: C \rightarrow B$ is another cocone, define a function $m: A \rightarrow B$ by

$$m(a) = f_{A\langle a_1 \rangle}(a_1) + if_{A\langle a_2 \rangle}(a_2).$$

This is a function satisfying $f_C = m \circ i_C$, and if it is a partial *-morphism then it is the unique such function, because a_1 and a_2 are uniquely defined by a . It remains to show that m is a well-defined morphism of **PCstar**, which follows from the assumption that the f_C are. Let us demonstrate the exemplaric clauses of preservation of scalar multiplication and multiplication. For $z \in \mathbb{C}$ and normal $a \in A$ we have $(za)_1 = xa_1 - ya_2$ and $(za)_2 = xa_2 + ya_1$, where $z = x + iy$ for $x, y \in \mathbb{R}$. Hence

$$\begin{aligned} m(za) &= f_{A\langle xa_1 - ya_2 \rangle}(xa_1 - ya_2) + if_{A\langle xa_2 + ya_1 \rangle}(xa_2 + ya_1) \\ &= f_{A\langle a_1, a_2 \rangle}(xa_1 - ya_2) + if_{A\langle a_1, a_2 \rangle}(xa_2 + ya_1) \\ &= xf_{A\langle a_1 \rangle}(a_1) + if_{A\langle a_2 \rangle}(a_2) + iyf_{A\langle a_1 \rangle}(a_1) - yf_{A\langle a_2 \rangle}(a_2) \\ &= (x + iy)(f_{A\langle a_1 \rangle}(a_1) + if_{A\langle a_2 \rangle}(a_2)) \\ &= zm(a). \end{aligned}$$

For commensurable $a, b \in A$, we have $(ab)_1 = a_1b_1 - a_2b_2$ and $(ab)_2 = a_1b_2 + a_2b_1$, so that

$$\begin{aligned} m(ab) &= f_{A\langle a_1b_1 - a_2b_2 \rangle}(a_1b_1 - a_2b_2) + if_{A\langle a_1b_2 + a_2b_1 \rangle}(a_1b_2 + a_2b_1) \\ &= f_{A\langle a_1b_1 - a_2b_2 \rangle}(a_1b_1) - f_{A\langle a_1b_1 - a_2b_2 \rangle}(a_2b_2) \\ &\quad + if_{A\langle a_1b_2 - a_2b_1 \rangle}(a_1b_2) + if_{A\langle a_1b_2 - a_2b_1 \rangle}(a_2b_1) \\ &= f_{A\langle a_1, b_1, a_2, b_2 \rangle}(a_1b_1) - f_{A\langle a_1, b_1, a_2, b_2 \rangle}(a_2b_2) \\ &\quad + if_{A\langle a_1, b_1, a_2, b_2 \rangle}(a_1b_2) + if_{A\langle a_1, b_1, a_2, b_2 \rangle}(a_2b_1) \\ &= f_{A\langle a_1, b_1 \rangle}(a_1b_1) + if_{A\langle a_1, b_2 \rangle}(a_1b_2) + if_{A\langle a_2, b_1 \rangle}(a_2b_1) - f_{A\langle a_2, b_2 \rangle}(a_2b_2) \\ &= f_{A\langle a_1 \rangle}(a_1)f_{A\langle b_1 \rangle}(b_1) + if_{A\langle a_1 \rangle}(a_1)f_{A\langle b_2 \rangle}(b_2) \\ &\quad + if_{A\langle a_2 \rangle}(a_2)f_{A\langle b_1 \rangle}(b_1) - f_{A\langle a_2 \rangle}(a_2)f_{A\langle b_2 \rangle}(b_2) \\ &= (f_{A\langle a_1 \rangle}(a_1) + if_{A\langle a_2 \rangle}(a_2))(f_{A\langle b_1 \rangle}(b_1) + if_{A\langle b_2 \rangle}(b_2)) \\ &= m(a)m(b). \end{aligned} \quad \square$$

Together, Theorem 15 above and Theorem 16 below show that, up to partial *-isomorphism, every C*-algebra can be reconstructed from its commutative C*-subalgebras, lending force to the Bohrification programme. In this light Theorem 15 could be said to embody a categorical crude version of complementarity.

Theorem 15 would fail, however, if we would drop the requirement that \odot is reflexive. The reason is the following: there exists a von Neumann algebra A

that is not anti-isomorphic to itself [2]; since this A is in standard form, *i.e.* has a separating cyclic vector, it follows that A is not isomorphic to its commutant A' . But Tomita-Takesaki theory shows that any von Neumann algebra A is anti-isomorphic to its commutant A' , whence $\mathcal{C}(A) \cong \mathcal{C}(A')$. Therefore not every C^* -algebra is the colimit of its commutative subalgebras in the category **CStar**. In addition, this result, combined with Theorem 15, provides a partial $*$ -isomorphism $N(A) \rightarrow N(A')$ that cannot be of the form $N(f)$ for some $f: A \rightarrow A'$. To put it another way, the faithful functor N is not full.

We close this section with a discussion of completeness and cocompleteness properties of **PCstar**. It is known that the category of C^* -algebras is both complete and cocomplete (for coproducts, see [16], and for coequalizers, see [8]). As it turns out, also **PCstar** is both complete and cocomplete.

PCstar is complete, as it has both equalizers and arbitrary products. Equalizers of partial C^* -algebras are constructed as in **Set**, having inherited commensurability and subalgebra structure. Products are given by $\prod_i A_i = \{(a_i)_i \mid a_i \in A_i, \sup_i \|a_i\| < \infty\}$, with componentwise commensurability and algebraic structure. In particular, the terminal object **1** is the 0-dimensional (partial) C^* -algebra $\{0\}$, confusingly sometimes also denoted by 0.

The coproduct of a family A_i of partial C^* -algebras is got by taking their disjoint union, identifying for every $z \in \mathbb{C}$ the elements of the form $z1_i$. Notice that elements from different summands A_i are never commensurable in the coproduct. In particular, the initial object **0** is the 1-dimensional (partial) C^* -algebra \mathbb{C} , which is confusingly sometimes also denoted by 1.

Coequalizers are harder to describe constructively, but they do exist.

Theorem 16 *The category **PCstar** is complete and cocomplete.*

PROOF To show that **PCstar** has coequalizers, the same strategy as the proof of Theorem 5 applies, because for every partial C^* -algebra A the collection of isomorphism classes of partial $*$ -maps $f: A \rightarrow B$ such that $f(B)$ is dense in A form a set and every partial C^* -algebra map with domain A factors through a map of this form. \square

4.1 Variations

Theorem 15 holds for many varieties of (partial) C^* -algebras, as its proof only depends on (partial) algebraic properties. Let us consider (partial) Rickart C^* -algebras as an example. Recall that a commutative C^* -algebra A is Rickart when every $a \in A$ has a unique projection $\text{RP}(a) \in A$ such that $(1 - \text{RP}(a)) \cdot A$ is the right annihilator $\{b \in A \mid ab = 0\}$. We call a partial C^* -algebra A together with a total map

$$\text{RP}: A \rightarrow A$$

a *partial Rickart C^* -algebra* when every pairwise commensurable $S \subseteq A$ is contained in a pairwise commensurable $T \subseteq A$ on which the operations of A yield a commutative Rickart C^* -algebra structure with RP's given by the function above. Denote the subcategory of **PCstar** whose objects are partial Rickart

C*-algebras and whose morphism are partial *-morphisms which preserve RP by **PRCstar**.

Theorem 17 *Every partial Rickart C*-algebra is the colimit of its commutative Rickart C*-subalgebras.*

PROOF The proof of Theorem 15 holds verbatim when every reference to (partial) C*-algebras is replaced by (partial) Rickart C*-algebras. \square

If **RCstar** is the subcategory of **Cstar** consisting of Rickart C*-algebras and *-morphisms preserving RP, then there is a functor

$$N : \mathbf{RCstar} \rightarrow \mathbf{PRCstar}$$

sending every Rickart C*-algebra A to its normal part; this follows from [1, Proposition 4.4]. Similar results hold for any type of C*-algebra that is defined by algebraic properties, such as AW*-algebras and spectral C*-algebras (see [9, 5.1]). We will come back to AW*-algebras in Section 6 below.

The distinguishing feature of von Neumann algebras amongst C*-algebras, in contrast, is topological in nature. This makes it harder to come up with a notion of partial von Neumann algebra: the obvious definition – a partial C*-algebra A in which every subset of commensurable elements is contained in a von Neumann algebra – has the drawback that it is not clear if $N(A)$ would be a partial von Neumann algebra given a von Neumann algebra A . We can, however, still obtain the following.

Let A be a von Neumann algebra. Without loss of generality, we may assume that A acts on a Hilbert space H . Denote the von Neumann subalgebra of a von Neumann algebra A generated by a subset $S \subseteq A$ by $A\langle\langle S \rangle\rangle$. It is the closure of the C*-algebra $A\langle S \rangle$ in the weak operator topology, and by von Neumann's double commutant theorem [12, Theorem 5.3.1], it equals $A\langle S \rangle''$.

Lemma 18 *If a C*-subalgebra C of a von Neumann algebra A is commutative, then so is its von Neumann envelope $A\langle\langle C \rangle\rangle$. Hence if $a \in A$ is normal, then $A\langle\langle a \rangle\rangle$ is commutative.*

PROOF Let $a, b \in C''$. Since C'' is the (weak operator) closure of C , we can write b as a (weak operator) limit $b = \lim_n b_n$ for $b_n \in C$. Then:

$$\begin{aligned} ab &= a(\lim_n b_n) = \lim_n ab_n && \text{(by [12, 5.7.9(i)])} \\ &= \lim_n b_n a && \text{(since } a \in C'' \text{ and } b_n \in C \subseteq C') \\ &= (\lim_n b_n)a && \text{(by [12, 5.7.9(ii)])} \\ &= ba. && \square \end{aligned}$$

Theorem 19 *Let A be a von Neumann algebra acting on a Hilbert space. Then $N(A)$ is a colimit in **PCstar** of the (finitely generated) commutative von Neumann subalgebras of A .*

PROOF Using Lemma 18, the proof of Theorem 15 holds verbatim when every occurrence of $A\langle S \rangle$ is replaced by $A\langle\langle S \rangle\rangle$. \square

5 Gelfand duality

The full subcategory of **PCstar** consisting of commutative C^* -algebras is just the category **cCstar** of commutative C^* -algebras and $*$ -morphisms. This category is dual to the category of compact Hausdorff spaces and continuous functions via Gelfand duality [10]. Constructively, the latter category is replaced by that of compact completely regular locales [3]:

$$\mathbf{cCstar} \begin{array}{c} \xrightarrow{\Sigma} \\ \xleftarrow[\mathbf{Loc}(-, \mathbb{C})]{\sim} \end{array} \mathbf{KRegLoc}^{\text{op}}, \quad (2)$$

where $\Sigma(A)$ is the Gelfand spectrum of a commutative C^* -algebra A . The dualizing object \mathbb{C} is both a locale and a (partial) C^* -algebra; recall that it is in fact the initial partial C^* -algebra $\mathbf{0}$.

The colimit theorem, Theorem 15, together with the fact that the categories in (2) are cocomplete and complete, enables us to prove the following extension of Gelfand duality.

Proposition 20 *There is a reflection*

$$\mathbf{PCstar} \begin{array}{c} \xrightarrow{K} \\ \xleftarrow[\mathbf{Loc}(-, \mathbb{C})]{\perp} \end{array} \mathbf{KRegLoc}^{\text{op}},$$

in which the functor K is determined by $K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C)$.

PROOF Let A be a partial C^* -algebra and X a compact completely regular locale. Then there are bijective correspondences:

$$\begin{array}{ll} \underline{\underline{f: K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C) \rightarrow X}} & \text{(in } \mathbf{KRegLoc}^{\text{op}}) \\ \underline{\underline{\forall C \in \mathcal{C}(A). f_C: \Sigma(C) \rightarrow X}} & \text{(in } \mathbf{KRegLoc}^{\text{op}}) \\ \underline{\underline{\forall C \in \mathcal{C}(A). g_C: C \rightarrow \mathbf{Loc}(X, \mathbb{C})}} & \text{(in } \mathbf{cCstar}) \\ \underline{\underline{g: A \rightarrow \mathbf{Loc}(X, \mathbb{C})}} & \text{(in } \mathbf{PCstar}). \end{array}$$

The first correspondence holds by definition of limit, the middle correspondence holds by Gelfand duality (2), and the last correspondence holds by Theorem 15. Since all correspondences are natural in A and X , this establishes the adjunction $K \dashv \mathbf{Loc}(-, \mathbb{C})$. Finally, since a commutative C^* -algebra is trivially a colimit of itself in **PCstar**, the adjunction is a reflection. \square

Theorem 21 *The reflection $K \dashv \mathbf{Loc}(-, \mathbb{C})$ extends Gelfand duality, i.e. the following diagram commutes (serially).*

$$\begin{array}{ccc} & \mathbf{cCstar} & \\ & \swarrow & \searrow \\ \mathbf{PCstar} & & \mathbf{KRegLoc}^{\text{op}} \\ & \xleftarrow[\mathbf{Loc}(-, \mathbb{C})]{\perp} & \\ & \xrightarrow{K} & \end{array}$$

$\mathbf{Loc}(-, \mathbb{C})$ (between \mathbf{cCstar} and $\mathbf{KRegLoc}^{\text{op}}$)
 Σ (between \mathbf{cCstar} and $\mathbf{KRegLoc}^{\text{op}}$)
 \sim (between \mathbf{cCstar} and $\mathbf{KRegLoc}^{\text{op}}$)

PROOF If A is a commutative C^* -algebra, it is the initial element in the diagram $\mathcal{C}(A)^{\text{op}}$ by Proposition 14(b). Hence $K(A) = \lim_{C \in \mathcal{C}(A)^{\text{op}}} \Sigma(C) = \Sigma(A)$. \square

Corollary 22 *Commutative C^* -algebras form a reflective full subcategory of partial C^* -algebras, i.e. the inclusion $\mathbf{cCstar} \hookrightarrow \mathbf{PCstar}$ has a left adjoint $L: \mathbf{PCstar} \rightarrow \mathbf{cCstar}$.*

PROOF The adjunctions of the previous theorem compose, giving the required left adjoint as $L = \mathbf{Loc}(-, \mathbb{C}) \circ K$. \square

This means that for a partial C^* -algebra A one has

$$\mathbf{PCstar}(A, \mathbb{C}) \cong \mathbf{cCstar}(L(A), \mathbb{C}).$$

In other words, multiplicative quasi-states of A which are multiplicative on commutative subalgebras precisely correspond to states of $L(A)$. Thus these quasi-states have good (categorical) behaviour. However, things are not as interesting as they may seem. By the Kochen-Specker theorem, no von Neumann algebra A without factors of type I_1 or I_2 can have such states ([4], see also [17]). It follows that $K(A) = \mathbf{0}$ and hence $L(A) = \mathbf{1}$ for such algebras. More generally, let us call a partial C^* -algebra A *Kochen-Specker* when $L(A) = \mathbf{1}$. Any such algebra A has no quasi-states: $\mathbf{PCstar}(A, \mathbb{C}) \cong \mathbf{cCstar}(\mathbf{1}, \mathbb{C}) = \emptyset$. Also, Kochen-Specker partial C^* -algebras are a ‘coproduct-ideal’ in a sense that we now make precise. For $X \in \mathbf{KRegLoc}$ we have $\mathbf{0} \times X = \mathbf{0}$, so by Gelfand duality (2), we have $\mathbf{1} + C = \mathbf{1}$ for a commutative C^* -algebra C . So if $A \in \mathbf{PCstar}$ is Kochen-Specker, and $B \in \mathbf{PCstar}$ arbitrary, then also $A + B$ is Kochen-Specker:

$$L(A + B) = L(A) + L(B) = \mathbf{1} + L(B) = \mathbf{1}.$$

The first equality holds because L , being a left adjoint, preserves colimits.

6 Projections, partial AW*-algebras and tensor products

This section discusses a functor $\mathbf{PCstar} \rightarrow \mathbf{PBoolean}$, relating Sections 2 and 3 to Sections 4 and 5.

6.1 Projections and partial AW*-algebras

An element p of a partial C^* -algebra A is called a *projection* when it satisfies $p^* = p = p^2$. The elements $0 \in A$ and $1 \in A$ are trivially projections; other projections are called nontrivial.

Lemma 23 *There is a functor $\text{Proj}: \mathbf{PCstar} \rightarrow \mathbf{PBoolean}$ where $\text{Proj}(A)$ is the set of projections of A .*

PROOF First, $\text{Proj}(A)$ is indeed a partial Boolean algebra. Commeasurability is inherited from A . One easily checks that $\neg p = 1 - p$ is a projection when p is. If p, q are commeasurable in $\text{Proj}(A)$, then they commute, whence the projection $p \wedge q = pq$ is also in A [17, 4.14]. This makes $\text{Proj}(A)$ into a partial Boolean algebra. Finally, morphisms of partial C^* -algebras are easily seen to preserve projections, making the assignment $A \mapsto \text{Proj}(A)$ functorial. \square

For the following class of partial C^* -algebras we get stronger results.

Definition 24 A partial Rickart C^* -algebra A is a *partial AW^* -algebra*, if it comes equipped with an operation

$$\bigvee: \{X \subseteq \text{Proj}(A) \mid X \times X \subseteq \odot\} \rightarrow \text{Proj}(A),$$

in such a way that each pairwise commeasurable $S \subseteq A$ is contained in a pairwise commeasurable $T \subseteq A$ on which the operations determine a commutative AW^* -algebra structure (*i.e.* the structure is that of a commutative Rickart C^* -algebra, whose projections form a complete Boolean algebra with suprema given by the operation \bigvee above). Denote the subcategory of **PCstar** whose objects are partial AW^* -algebras and whose morphisms are partial $*$ -morphisms which preserve RP and \bigvee by **PAWstar**.

Lemma 25 *The functor Proj restricts to a functor $\mathbf{PAWstar} \rightarrow \mathbf{PCBoolean}$.*

PROOF Clear from the definition of a partial AW^* -algebra. \square

Remark 26 It would be interesting to see whether this functor is part of an equivalence, like in the total case, where it is one side of an equivalence of categories between **cAWstar** and **CBoolean**.

Proposition 27 *The functors Proj and \mathcal{C} commute for partial AW^* -algebras: writing \mathcal{C}' for the functor $\mathbf{PAWstar} \rightarrow [\mathbf{POrder}, \mathbf{cAWstar}]$, and \mathcal{C} for the functor $\mathbf{PBoolean} \rightarrow \mathbf{POrder}$, we have $\mathcal{C} \circ \text{Proj} = \text{Proj} \circ \mathcal{C}'$. Explicitly,*

$$\{\text{Proj}(C) \mid C \in \mathcal{C}(A)\} = \mathcal{C}(\text{Proj}(A))$$

for every partial AW^* -algebra A .

PROOF This follows from the combination of Stone and Gelfand duality, which yields an equivalence between **cAWstar** and **CBoolean**. One direction of the equivalence is obtained by taking projections and the other is obtained by taking $C(X)$ where X is the Stone space associated to the (complete) Boolean algebra. For the purposes of the proof, we will denote the latter composite functor by F . In particular, the projections $\text{Proj}(C)$ of a commutative AW^* -algebra C form a complete Boolean algebra and the left-hand side is contained in the right-hand side.

For the converse, let B be a complete Boolean lattice of projections in A . Then the projections in B commute pairwise, and hence generate a commutative

AW*-subalgebra $C = A\langle B \rangle$. We obviously have an inclusion $i : B \subseteq \text{Proj}(C)$. But then the composite

$$FB \xrightarrow{Fi} F\text{Proj}(C) \xrightarrow[\cong]{\eta_C^{-1}} C,$$

where η is the unit of the adjunction $\text{Proj} \dashv F$, shows that FB is isomorphic to a commutative AW*-subalgebra of A contained in C . This commutative subalgebra also contains B , because the diagram

$$\begin{array}{ccc} \text{Proj}(FB) & \xrightarrow{\text{Proj}(Fi)} & \text{Proj}(F\text{Proj}(C)) \\ \epsilon_B \downarrow \cong & & \cong \downarrow \epsilon_{\text{Proj}(C)} = \text{Proj}(\eta_C^{-1}) \\ B & \xrightarrow{i} & \text{Proj}(C) \end{array}$$

commutes by naturality of the counit $\epsilon : \text{Proj}(F-) \Rightarrow 1$. Since $C = A\langle B \rangle$, this implies that Fi is an isomorphism. But then so is i and therefore $B = \text{Proj}(C)$. \square

As a corollary to the previous proposition, we can extend Proposition 14 with atoms to mirror Proposition 2. Keep in mind that the following corollary does not entail that $\mathcal{C}(A)$ is atomic.

Corollary 28 *For a partial AW*-algebra A , the atoms of the poset $\mathcal{C}(A)$ are $A\langle p \rangle$ for nontrivial projections p . \square*

6.2 Tensor products

It is clear from the description of coproducts in **PCstar** and **PBoolean** that the functor Proj preserves coproducts. Recall that in a coproduct of partial Boolean or C*-algebras, nontrivial elements from different summands are never commensurable. Theorems 4 and 15 provide the option of defining a tensor product satisfying the adverse universal property.

Definition 29 Let A and B be a pair of partial Boolean algebras (partial C*-algebras). Define

$$A \otimes B = \text{colim}\{C + D \mid C \in \mathcal{C}(A), D \in \mathcal{C}(B)\},$$

where $C + D$ is the coproduct in the category of Boolean algebras (commutative C*-algebras).

There are canonical morphisms $\kappa_A : A \rightarrow A \otimes B$ and $\kappa_B : B \rightarrow A \otimes B$ as follows. By definition, $A \otimes B$ is the colimit of $C + D$ for $C \in \mathcal{C}(A)$ and $D \in \mathcal{C}(B)$. Precomposing with the coproduct injections $C \rightarrow C + D$ gives a cocone $C \rightarrow A \otimes B$ on $\mathcal{C}(A)$. By the colimit theorem, A is the colimit of $\mathcal{C}(A)$. Hence there is a mediating morphism $\kappa_A : A \rightarrow A \otimes B$.

The unit element for both the tensor product and the coproduct is the initial object $\mathbf{0}$. The big difference between $A \otimes B$ and the coproduct $A + B$ is that elements $\kappa_A(a)$ and $\kappa_B(b)$ are always commensurable in the former, but never in the latter. Indeed, this universal property characterizes the tensor product.

Proposition 30 *Let $f: A \rightarrow Z$ and $g: B \rightarrow Z$ be morphisms in the category **PBoolean** (**PCstar**). The cotuple $[f, g]: A + B \rightarrow Z$ factorizes through $A \otimes B$ if and only if $f(a) \odot f(b)$ for all $a \in A$ and $b \in B$.*

PROOF By construction, giving $h: A \otimes B \rightarrow Z$ amounts to giving a cocone $C + D \rightarrow Z$ for $C \in \mathcal{C}(A)$ and $D \in \mathcal{C}(B)$. Because $C + D$ is totally defined, any morphism $C + D \rightarrow Z$ must also be total. But this holds (for all C and D) if and only if $f(a)$ and $g(b)$ are commeasureable for all $a \in A$ and $b \in B$, for (only) then can one take h to be the cotuple of the corestrictions of f and g . \square

The tensor products of Definition 29 makes $\text{Proj}: \mathbf{PCstar} \rightarrow \mathbf{PBoolean}$ a monoidal functor: the natural transformation $\text{Proj}(A) \otimes \text{Proj}(B) \rightarrow \text{Proj}(A \otimes B)$ is induced by the cotuples $\text{Proj}(C) + \text{Proj}(D) \rightarrow \text{Proj}(A \otimes B)$ of

$$\text{Proj}(C) \xrightarrow{\text{Proj}(C \hookrightarrow A)} \text{Proj}(A) \xrightarrow{\text{Proj}(\kappa_A)} \text{Proj}(A \otimes B).$$

Proposition 31 *The functor $\text{Proj}: \mathbf{PCstar} \rightarrow \mathbf{PBoolean}$ preserves coproducts and is monoidal.* \square

We end this section by discussing the relation between the tensor products of partial Boolean algebras and those of Hilbert spaces, describing compound quantum systems. Let **Hilb** be the category of Hilbert spaces and continuous linear maps, and let $B: \mathbf{Hilb} \rightarrow \mathbf{PCstar}$ denote the functor $B(H) = \mathbf{Hilb}(H, H)$ acting on morphisms as $B(f) = f \circ (-) \circ f^\dagger$. The definition of the tensor product in **PCstar** as a colimit yields a natural transformation $B(H) \otimes B(K) \rightarrow B(H \otimes K)$, induced by morphisms $C \rightarrow B(H \otimes K)$ for $C \in \mathcal{C}(B(H))$ given by $a \mapsto a \otimes \text{id}_K$. Initiality of the tensor unit $\mathbf{0}$ gives a morphism $\mathbf{0} \rightarrow B(\mathbb{C})$, and these data satisfy the coherence requirements. Hence the functor B is monoidal, and therefore also the composite $\text{Proj} \circ B: \mathbf{Hilb} \rightarrow \mathbf{PBoolean}$ is a monoidal functor.

7 Functoriality of Bohrfication

The so-called Bohrfication construction (see [9], whose notation we adopt) associates to every C*-algebra A an internal commutative C*-algebra \underline{A} in the topos $[\mathcal{C}(A), \mathbf{Set}]$, given by the tautological functor $\underline{A}(C) = C$. Gelfand duality then yields an internal locale, which can in turn be externalized. As it happens this construction works equally well for partial C*-algebras, so that Bohrfication for ordinary C*-algebras can be seen as the composition of the functor N from Proposition 13 with Bohrfication for partial C*-algebras. Thus a locale is associated to every object of **PCstar**. In this final section we consider its functorial aspects. It turns out that the whole construction summarized above can be made into a functor from partial C*-algebras to locales by restricting the morphisms of the former.

Bohrfication does not just assign a topos to each (partial) C*-algebra, it assigns a topos with an internal C*-algebra. To reflect this, we define categories of toposes equipped with internal structures.

Definition 32 The category **RingedTopos** has as objects pairs (T, R) of a topos T and an internal ring object $R \in T$. A morphism $(T, R) \rightarrow (T', R')$ consists of a geometric morphism $F: T' \rightarrow T$ and an internal ring morphism $\varphi: R' \rightarrow F^*(R)$ in T .

By **CstaredTopos** we denote the subcategory of **RingedTopos** of objects (T, A) where A is an internal C^* -algebra in T and morphisms (F, φ) where φ is an internal $*$ -ring morphism.

Notice that the direction of morphisms in this definition is opposite to the customary one in algebraic geometry [7, 4.1].

First of all, any functor $\mathbf{D} \rightarrow \mathbf{C}$ induces a geometric morphism $[\mathbf{D}, \mathbf{Set}] \rightarrow [\mathbf{C}, \mathbf{Set}]$, of which the inverse part is given by precomposition (see [11, A4.1.4]). We have already seen that \mathcal{C} is a functor $\mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{POrder}^{\text{op}}$. Additionally, restricting a morphism $f: B \rightarrow A$ of partial C^* -algebras to $D \in \mathcal{C}(B)$ and corestricting to $\mathcal{C}f(D)$ gives a morphism of commutative C^* -algebras. Hence we obtain a geometric morphism of toposes $[\mathcal{C}(B), \mathbf{Set}] \rightarrow [\mathcal{C}(A), \mathbf{Set}]$ as well as an internal morphism of commutative $*$ -rings. The latter is a natural transformation whose component at D is $\underline{B}(D) \rightarrow ((\mathcal{C}f)^* \underline{A})(D) = \underline{A}(\mathcal{C}f(D))$. In other words, we have a functor $\mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{RingedTopos}$.

In general, (inverse parts of) geometric morphisms do not preserve internal C^* -algebras. But in this particular case, $(\mathcal{C}f)^* \underline{A}$ is in fact an internal C^* -algebra in $[\mathcal{C}(B), \mathbf{Set}]$. The proof is contained in [9, 4.8], which essentially shows that any functor from a poset P to the category of C^* -algebras is always an internal C^* -algebra in the topos $[P, \mathbf{Set}]$. Therefore, we really have a functor $\mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{CstaredTopos}$, as the following proposition records.

Proposition 33 *Bohrification is functorial* $\mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{CstaredTopos}$. \square

Applying internal Gelfand duality to $\varphi: \underline{B} \rightarrow (\mathcal{C}f)^* \underline{A}$ gives an internal locale morphism $\underline{\Sigma}((\mathcal{C}f)^* \underline{A}) \rightarrow \underline{\Sigma}(\underline{B})$. But to get a functor $\mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{LocaledTopos}$, for the evident definition of the latter category, we would need an internal locale morphism $(\mathcal{C}f)^*(\underline{\Sigma}(\underline{A})) \rightarrow \underline{\Sigma}(\underline{B})$. As $(\mathcal{C}f)^*(\underline{\Sigma}(\underline{A}))$ and $\underline{\Sigma}((\mathcal{C}f)^* \underline{A})$ are incomparable in general, this is where the current line of reasoning stops. The most natural way out is to restrict the morphisms of \mathbf{PCstar} as follows.

Lemma 34 *For morphisms $f: A \rightarrow B$ of \mathbf{PCstar} , the following are equivalent:*

- (a) *if $\mathcal{C}f(C) \leq D$ and $\mathcal{C}f(C') \leq D$ for $C, C' \in \mathcal{C}(A)$ and $D \in \mathcal{C}(B)$, then there is $C'' \in \mathcal{C}(A)$ such that $C \leq C''$ and $C' \leq C''$ and $\mathcal{C}f(C'') \leq D$;*
- (b) *$a \odot a'$ when $f(a) \odot f(a')$.*

PROOF First assume (a) and suppose $f(a) \odot f(a')$. Take $C = A\langle a, a^* \rangle$, $C' = A\langle a', (a')^* \rangle$, and $D = B\langle f(a), f(a'), f(a)^*, f(a')^* \rangle$. Then $\mathcal{C}f(C) \leq D$ and $\mathcal{C}f(C') \leq D$. Hence there is C'' with $C \leq C''$ and $C' \leq C''$. So a, a' are both elements of the commutative algebra C'' , so $a \odot a'$.

Conversely, assuming (b) and supposing $\mathcal{C}f(C) \leq D$ and $\mathcal{C}f(C') \leq D$, for all $a \in C$ and $a' \in C'$ we have $f(a), f(a') \in D$, so that $f(a) \odot f(a')$. But that means that C and C' are commuting commutative subalgebras of A . Hence we can take $C'' = A\langle C, C' \rangle$. \square

We say that morphisms satisfying the conditions in the previous lemma *reflect commensurability*. Notice that this class of morphisms excludes the type of counterexample discussed after Theorem 4. To show how the assignment of a locale to a partial C*-algebra becomes functorial with these morphisms, let us switch to its external description [9, 5.16]:

$$S(A) = \{F: \mathcal{C}(A) \rightarrow \mathbf{Set} \mid F(C) \text{ open in } \Sigma(C), F \text{ monotone}\}. \quad (3)$$

For A a partial C*-algebra, $S(A)$ is a locale. We want to extend this to a functor $S: \mathbf{PCstar}^{\text{op}} \rightarrow \mathbf{Loc}$, or equivalently, a functor $S: \mathbf{PCstar} \rightarrow \mathbf{Frm}$. Let $f: A \rightarrow B$ be a morphism of partial C*-algebras, $F \in S(A)$, and $D \in \mathcal{C}(B)$. If $C \in \mathcal{C}(A)$ satisfies $\mathcal{C}f(C) \leq D$, then we have a morphism $C \rightarrow D$ given by the composition $C \xrightarrow{f} \mathcal{C}f(C) \leq D$. Its Gelfand transform is a frame morphism $\Sigma(C \xrightarrow{f} \mathcal{C}f(C) \leq D): \Sigma(C) \rightarrow \Sigma(D)$. So, since $F(C) \in \Sigma(C)$, we get an open in $\Sigma(D)$. The fact that $\Sigma(D)$ is a locale allows us to take the join over all such C , ending up with the candidate action on morphisms

$$Sf(F)(D) = \bigvee_{\substack{C \in \mathcal{C}(A) \\ \mathcal{C}f(C) \leq D}} \Sigma(C \rightarrow \mathcal{C}f(C) \leq D)(F(C)). \quad (4)$$

Theorem 35 *Bohrification gives a functor $S: \mathbf{PCstar}_{\text{rc}}^{\text{op}} \rightarrow \mathbf{Loc}$, where the domain is the subcategory of \mathbf{PCstar} of morphisms reflecting commensurability.*

PROOF One quickly verifies that $Sf(F)$, as given in (4), is monotone, and hence a well-defined element of $S(B)$, and that Sf preserves suprema. The greatest element $1 \in S(A)$ is also preserved by Sf :

$$Sf(1)(D) = \bigvee_{\substack{C \in \mathcal{C}(A) \\ \mathcal{C}f(C) \leq D}} \Sigma(C \rightarrow D)(1) = \bigvee_{\substack{C \in \mathcal{C}(A) \\ \mathcal{C}f(C) \leq D}} 1 = 1.$$

The last equality holds because the join is not taken over the empty set: there is always $C \in \mathcal{C}(A)$ with $\mathcal{C}f(C) \leq D$, namely $C = \mathbf{0}$.

To finish well-definedness and show that Sf is a frame morphism, we need to show that it preserves binary meets. Recall that any frame satisfies the infinitary distributive law $(\bigvee_i y_i) \wedge x = \bigvee_i (y_i \wedge x)$. It follows that one always has $(\bigvee_i y_i) \wedge (\bigvee_j x_j) \geq \bigvee_k (y_k \wedge x_k)$. A sufficient (but not necessary) condition for equality to hold would be if for all i, j there exists k such that $y_i \wedge x_j \leq y_k \wedge x_k$. Expanding the definition of Sf and writing $x_C = \Sigma(C \rightarrow D)(F(C))$

and $y_C = \Sigma(C \rightarrow D)(G(C))$ gives precisely this situation:

$$Sf(F \wedge G)(D) = \bigvee_{\mathcal{C}f(C'') \leq D} x_{C''} \wedge y_{C''},$$

$$(Sf(F) \wedge Sf(G))(D) = \left(\bigvee_{\mathcal{C}f(C) \leq D} x_C \right) \wedge \left(\bigvee_{\mathcal{C}f(C') \leq D} y_{C'} \right).$$

So, by Lemma 34, Sf will preserve binary meets if f reflects commeasureability. Finally, it is easy to see that $S(\text{id}) = \text{id}$ and $S(g \circ f) = Sg \circ Sf$. \square

We conclude with three remarks concerning the last theorem.

- Note that replacing the Gelfand spectrum by the Stone spectrum yields a similar functor $\mathbf{PBoolean}_{\text{rc}}^{\text{op}} \rightarrow \mathbf{Loc}$.
- The externalization of an internal locale in a topos $\text{Sh}(L)$ consists of a locale morphism $L' \rightarrow L$, and not just the locale L' itself. In that spirit, the previous theorem should have taken into account the map $S(A) \rightarrow \mathcal{C}(A)$ as well. To consider functorial dependence on A of this map would require to change the category \mathbf{Loc} into something more complicated, from which we refrain here.
- The category \mathbf{Loc} is \mathbf{POrder} -enriched, and hence a 2-category. We remark that the functor S given by (3) and (4) shows that the externalization of Bohrification is a two-dimensional colimit (in \mathbf{Loc}) of the Gelfand spectra of commeasureable subalgebras. So, interestingly, whereas a one-dimensional colimit of the spectra must be trivial because of the Kochen-Specker theorem, Bohrification exhibits that a two-dimensional colimit can be nontrivial.

Acknowledgement

We thank Klaas Landsman for useful comments. Additionally, we owe the observation concerning [2] to Thierry Coquand, and the reference to [6] to John Harding and Mirko Navara and their forthcoming paper ‘Subalgebras of orthomodular lattices’.

References

- [1] Sterling K. Berberian. *Baer rings and Baer *-rings*. Online book, 2003. Available at http://www.ma.utexas.edu/mp_arc/c/03/03-181.pdf.
- [2] Alain Connes. A factor not anti-isomorphic to itself. *Annals of Mathematics*, 101(3):536–554, 1975.
- [3] Thierry Coquand and Bas Spitters. Constructive Gelfand duality for C^* -algebras. *Mathematical Proceedings of the Cambridge Philosophical Society*, 147(2):323–337, 2009.

- [4] Andreas Döring. Kochen-Specker theorem for Von Neumann algebras. *International Journal of Theoretical Physics*, 44(2):139–160, 2005.
- [5] Andreas Döring and Christopher J. Isham. ‘What is a thing?’: Topos theory in the foundations of physics. In *New Structures for Physics*, Lecture Notes in Physics. Springer, to appear, 2009. arXiv:0803.0417.
- [6] G. Grätzer, K. M. Koh, and M. Makkai. On the lattice of subalgebras of a boolean algebra. *Proceedings of the American Mathematical Society*, 36:87–92, 1972.
- [7] Alexandre Grothendieck. Le langage des schémas. In *Éléments de géométrie algébrique*, volume 4, pages 5–228. Publications Mathématiques de l’IHÉS, 1960.
- [8] Alain Guichardet. Sur la catégorie des algèbres de Von Neumann. *Bulletin des sciences mathématiques, 2e série*, 90:41–64, 1966.
- [9] Chris Heunen, Nicolaas P. Landsman, and Bas Spitters. Bohrification. In *Deep Beauty*. Cambridge University Press, to appear, 2009. arXiv:0909.3468.
- [10] Peter T. Johnstone. *Stone spaces*. Number 3 in Cambridge studies in advanced mathematics. Cambridge University Press, 1982.
- [11] Peter T. Johnstone. *Sketches of an elephant: A topos theory compendium*. Oxford University Press, 2002.
- [12] Richard V. Kadison and John R. Ringrose. *Fundamentals of the theory of operator algebras*. Academic Press, 1983.
- [13] Gudrun Kalmbach. *Orthomodular Lattices*. Academic Press, 1983.
- [14] Simon Kochen and Ernst Specker. The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics*, 17:59–87, 1967.
- [15] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer, 2nd edition, 1971.
- [16] Daniel Rainjonneau. Existence des sommes dans certaines catégories d’algèbres. *Les Comptes Rendus de l’Académie des sciences Paris, Série A*, 262:283–285, 1966.
- [17] Miklós Rédei. *Quantum Logic in Algebraic Approach*. Kluwer, 1998.