

Optimization 2

Lecturer: Hao Tang

Definition 1. The minimum of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is written as $\min_x f(x)$, and has the property that $\min_x f(x) \leq f(y)$ for any y .

Definition 2. The value x^* such that $f(x^*) = \min_x f(x)$ is called a minimizer.

Example 1. For the parabola $f(x) = x^2 + 4x - 1 = (x + 2)^2 - 5$, the minimum is -5 and the minimizer is $x = -2$.

Definition 3. A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex if for any $0 \leq \alpha \leq 1$, we have

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y) \quad (1)$$

for any x and y .

Definition 4. A function f is concave if $-f$ is convex.

Example 2. If f is convex, then

$$f(x) \geq f(y) + \nabla f(y)^\top (x - y) \quad (2)$$

for any x and y .

We can arrange the following

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y) \quad (3)$$

into

$$f(y) + \frac{f(y + \alpha(x - y)) - f(y)}{\alpha} \leq f(x). \quad (4)$$

Remember that this holds for any $0 \leq \alpha \leq 1$. In particular, if we take the limit,

$$f(y) + \lim_{\alpha \rightarrow 0} \frac{f(y + \alpha(x - y)) - f(y)}{\alpha} = f(y) + \nabla f(y)^\top (x - y) \leq f(x). \quad (5)$$

Definition 5. A matrix A is positive semidefinite if $v^\top A v \geq 0$ for all v , and is written as $A \succeq 0$.

Example 3. A function is convex if its Hessian is positive semidefinite.

The proof relies on mean-value theorem. It's not difficult, but is beyond the scope of this course.

Example 4. Show that the mean-squared error $\ell(y, \hat{y}) = (y - \hat{y})^2$ is convex in \hat{y} .

$$\frac{\partial^2}{\partial \hat{y}^2} \ell = 2 \geq 0. \quad (6)$$

Example 5. Show that the function

$$f(x) = x^\top \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} x \quad (7)$$

is convex.

The Hessian of f is $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$. For any $v = [v_1 \ v_2]^\top$, we have

$$[v_1 \ v_2] \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = [2v_1 \ 3v_2] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = 2v_1^2 + 3v_2^2 \geq 0 \quad (8)$$

The Hessian of f is positive semidefinite.

Example 6. Show that the Hessian of $f(x) = \|x\|_2^2$ is $2I$, and hence $\|x\|_2^2$ is convex in x .

$$\frac{\partial^2}{\partial x_i \partial x_j} f = 0 \quad \frac{\partial^2}{\partial x_i^2} f = 2 \quad (9)$$

Example 7. Show that if f is convex, then $g(x) = f(Ax + b)$ is also convex.

$$g(\alpha x + (1 - \alpha)y) = f(\alpha(Ax + b) + (1 - \alpha)(Ay + b)) \quad (10)$$

$$\leq \alpha f(Ax + b) + (1 - \alpha)f(Ay + b) = \alpha g(x) + (1 - \alpha)g(y) \quad (11)$$

Example 8. Show that if f_1, \dots, f_k are convex, then $f = \beta_1 f_1 + \dots + \beta_k f_k$ is also convex when $\beta_1, \dots, \beta_k \geq 0$.

$$f(\alpha x + (1 - \alpha)y) = \beta_1 f_1(\alpha x + (1 - \alpha)y) + \cdots + \beta_k f_k(\alpha x + (1 - \alpha)y) \quad (12)$$

$$\leq \beta_1 \alpha f_1(x) + \beta_1 (1 - \alpha) f_1(y) + \cdots + \beta_k \alpha f_k(x) + \beta_k (1 - \alpha) f_k(y) \quad (13)$$

$$= \alpha(\beta_1 f_1(x) + \cdots + \beta_k f_k(x)) + (1 - \alpha)(\beta_1 f_1(y) + \cdots + \beta_k f_k(y)) \quad (14)$$

$$= \alpha f(x) + (1 - \alpha) f(y) \quad (15)$$

Exercise 1. Given a data set of n samples $\{(x_1, y_1), \dots, (x_n, y_n)\}$, show that

$$L = \sum_{i=1}^n (w^\top x_i - y_i)^2 = \|Xw - y\|_2^2 \quad (16)$$

if we have

$$X = \begin{bmatrix} - & x_1 & - \\ & \vdots & \\ - & x_n & - \end{bmatrix} \quad y = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}. \quad (17)$$

Exercise 2. Given a data set of n samples $\{(x_1, y_1), \dots, (x_n, y_n)\}$, show that the mean-squared error

$$L = \|Xw - y\|_2^2 \quad (18)$$

is convex.

Example 9. Show that if f is convex and $\nabla f(x^*) = 0$ for a point x^* , then x^* is the minimizer of f .

Because f is convex, we have for any x and y ,

$$f(x) \geq f(y) + \nabla f(y)^\top (x - y). \quad (19)$$

In particular, if we let $y = x^*$,

$$f(x) \geq f(x^*) + \nabla f(x^*)^\top (x - x^*) = f(x^*). \quad (20)$$

Example 10. Show that $\nabla_x (x^\top A x) = (A^\top + A)x$.

We see that $x^\top A x$ is a real value. If we take the derivative of $x^\top A x$, we get

$$\frac{\partial}{\partial x_k} \sum_{i=1}^d \sum_{j=1}^d a_{ij} x_i x_j = \sum_{i \neq j}^d a_{ik} x_i + \sum_{j \neq i}^d a_{kj} x_j + \sum_{i=1}^d 2a_{ii} x_i \quad (21)$$

$$= \sum_{i=1}^d a_{ik} x_i + \sum_{j=1}^d a_{kj} x_j = a_{\cdot k}^\top x + a_{k \cdot} x \quad (22)$$

where $a_{\cdot k}$ is the k -th column of A and a_k is the k -th row of A .

Example 11. Show that $w^* = (X^\top X)^{-1} X^\top y$ is the minimizer for $L = \|Xw - y\|_2^2$.

$$L = (Xw - y)^\top (Xw - y) = w^\top X^\top Xw - 2y^\top Xw + y^\top y \quad (23)$$

$$\nabla L = (X^\top X + X^\top X)w - 2X^\top y = 0 \quad (24)$$

If $w^* = (X^\top X)^{-1} X^\top y$, then $\nabla L(w^*) = 0$. Because L is convex in w , w^* is a minimizer of L .

Example 12. Show that $\ell(s) = \log(1 + \exp(-s))$ is convex in s .

$$\frac{\partial \ell}{\partial s} = \frac{-\exp(-s)}{1 + \exp(-s)} = \frac{1}{1 + \exp(-s)} - 1 \quad (25)$$

$$\frac{\partial^2 \ell}{\partial s^2} = \frac{-1}{1 + \exp(-s)} \frac{-\exp(-s)}{1 + \exp(-s)} = \frac{1}{1 + \exp(-s)} \left(1 - \frac{1}{1 + \exp(-s)} \right) \geq 0 \quad (26)$$

Exercise 3. Given a data set of n samples $\{(x_1, y_1), \dots, (x_n, y_n)\}$, show that the log loss

$$L = \sum_{i=1}^n \log \left(1 + \exp(-y_i w^\top x_i) \right) \quad (27)$$

is convex.

Definition 6. A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is called strictly convex if for $0 \leq \alpha \leq 1$, we have

$$f(\alpha x + (1 - \alpha)y) < \alpha f(x) + (1 - \alpha)f(y) \quad (28)$$

for any $x \neq y$.

Exercise 4. A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is strictly convex if

$$f(x) > f(y) + \nabla f(y)^\top (x - y) \quad (29)$$

for any $x \neq y$.

Definition 7. A matrix A is positive definite if $v^\top A v > 0$ for any $v \neq 0$.

Exercise 5. A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is strictly convex if its Hessian is positive definite.

Example 13. Show that if f is strictly convex, then f has a unique minimizer.

Suppose x^* is a minimizer of f , i.e., $\nabla f(x^*) = 0$. The inequality

$$f(x) > f(y) + \nabla f(y)^\top (x - y). \quad (30)$$

holds for any $x \neq y$. In particular, if we let $y = x^*$,

$$f(x) > f(x^*) + \nabla f(x^*)^\top (x - x^*) = f(x^*). \quad (31)$$